

**SUPPLEMENTS**

**NEORNI  
PHARMA**



**2025**



# NeorniPharma Supplements

## Preventive formulations tailored to your birds

This catalogue is a comprehensive guide of the NeorniPharma range aimed at **disease prevention through natural, highly concentrated supplements**. This range offers a safe and successful alternative to medicinal treatments and a long-term solution. By targeting annually recurring problems, NeorniPharma aims to build immunity and consequently stronger birds.



### NeorniPharma Magistral Preparations

In addition to supplements, NeorniPharma also develops **Magistral preparations specifically aimed at disease control**. With this, NeorniPharma offers a short-term solution. Some of these products are featured in this catalogue. The full 'blue catalogue' is available from NeorniVet. Magistral preparations are only available on veterinary prescription.



**NEW**

#### New front desk address

Beverlosesteenweg 129  
3583 Paal, Belgium

#### New telephone number

+32 (0)11 96 91 02

#### Additional dosages of powders

View alternative dosages on our 'Powders and capsules' dosage card in this catalogue. View the full dosage card on page 16.



3 / 1

### Bird species in this catalogue



Pigeons



Songbirds



Fowl



Parrots



Birds of prey

Download our catalogues



All NeorniPharma products have unique formulas and are designed by veterinarians based on scientific research and practical experience.

## Breeding & immunity

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Immuno-Booster NEW (1 kg)	20 g/kg egg food/feed	All-in premix	65
Immuno-Plus (100 ml)	2 ml/l	Breeding, show/competition, extra energy	25
Immuno-Plus (250 ml)	2 ml/l	Breeding, show/competition, extra energy	55
Immuno-Plus (1 l)	2 ml/l	Breeding, show/competition, extra energy	179
Immuno Recup Caps (80 capsules)	1 capsule/0.5-1 Kg body weight	Immunity, recovery, energy	24
Super Start-Up (200 g)	5 g/kg feed	Start-up supplement for chicks, black spot	25
Vit E + Se (100 ml)	2 ml/l	Extra vitamin E & selenium	17
Vit E + Se (250 ml)	2 ml/l	Extra vitamin E & selenium	35

## Liver, muscles and condition

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Hepa-Plus (100 ml)	4 ml/l	Liver support	20
Hepa-Plus (250 ml)	4 ml/l	Liver support	36
Hepa-Plus (1 l)	4 ml/l	Liver support	96
Ready (100 ml)	5 ml/l	Form, condition & improved blood flow	22
Ready (250 ml)	5 ml/l	Form, condition & improved blood flow	48
Ready (1 l)	5 ml/l	Form, condition & improved blood flow	144

## Megabacteria and dysbacteriosis

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Dysbac Liquid (250 ml)	2 ml/l	Crop, proventriculus & intestinal health	37

## Natural oils

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Feather Energy (400 ml)	5 ml/kg feed	Flight Oil	19
Freeson Drops (30 ml)	1-2 drops/naris	Unobstructed respiratory tract	17
PikStop (100 ml)	1 ml/l	Pecking	28
Respicor (100 ml)	3 ml/l	Respiratory tract, heat stress	19
Respicor (250 ml)	3 ml/l	Respiratory tract, heat stress	35

## Parasites

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Derma-Plus (50 ml)	8 ml/l/40 m <sup>2</sup>	Red mite, northern fowl mite	35
Exmite Spray (250 ml)	Spray	Ectoparasites	22
Exmite Spray (1 l)	Spray	Ectoparasites	69

## Probiotics and disinfection

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Neorni Stabilizer (1 l)	5-10 ml/100 ml	Probiotics – ambient	20
Nositol (50 ml)	1-2 drops/naris	Nasal cleanser	10
Probi-Plus (200 g)	3 g/l	Probiotics and prebiotics	18
SilverSpray (500 ml)	Spray	Disinfection spray	15
Skin Spray (200 ml)	Spray	Probiotics - skin infections	17

All advertised prices include VAT applicable in Belgium (21%)



## Trichomoniasis, candida, aspergillosis & drinking water disinfection

PRODUCT	DOSAGE	INDICATION	PRICE (€)
CU-For Liquid (250 ml)	1-2 ml/l	Drinking water acidifier	15
SilverFloX (1 l)	2 ml/l	Drinking water disinfectant	32
Trichoryza Drops (50 ml)	1 drop/naris	Throat elixir	17
Trichoryza Drops (250 ml)	10 ml/kg	Throat elixir	59
Trichoryza Drops (1 l)	10 ml/kg	Throat elixir	165

## Vitamins, amino acids, electrolytes, minerals & trace elements

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Aminovital (100 ml)	2 ml/l	Amino acids, electrolytes, vitamins	19
Aminovital (250 ml)	2 ml/l	Amino acids, electrolytes, vitamins	35
Aminovital (1 l)	2 ml/l	Amino acids, electrolytes, vitamins	65
Condi-Plus (100 ml)	2 ml/l	Moulting supplement	15
Condi-Plus (250 ml)	2 ml/l	Moulting supplement	26
Condi-Plus (1 l)	2 ml/l	Moulting supplement	83
Vita Energy (400 g)	10 g/kg feed	Condition powder - vitamins, minerals and amino acids	30
Vita-Min-DCP (100 ml) 2 ml/l		Vit D, calcium, phosphorus & other minerals	17
Vita-Min-DCP (250 ml) 2 ml/l		Vit D, calcium, phosphorus & other minerals	32
Vita-Min-DCP (1 l)	2 ml/l	Vit D, calcium, phosphorus & other minerals	88
Vita-Plus (100 ml)	2 ml/l	Water-soluble multivitamins	18
Vita-Plus (250 ml)	2 ml/l	Water-soluble multivitamins	28
Vita-Plus (1 l)	2 ml/l	Water-soluble multivitamins	65

## PURE

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Brewer's yeast (1 kg)	15-30 g/kg feed	Amino acids, minerals, trace elements, vitamin B	18
Dextrose (1 kg)	15-30 g/l	Energy, flavour	18
Electrolyte (500 g)	15-30 g/l	Hydration, stabilisation, balance	18
Yellow (100 g)	1-2 g/kg feed	Colour	15
Yellow (500 g)	1-2 g/kg feed	Colour	60
Highly digestible protein (400 g)	5-15 g/kg feed	Muscle growth, additional amino acids/proteins	25
Red (100 g)	Max. 1 g/kg feed	Colour	28
Red (500 g)	Max. 1 g/kg feed	Colour	85

# NEORNI

NeorniPharma is part of Neorni, a team of veterinarians dedicated specifically to avian medicine. Neorni offers various services to ensure the best care for birds.

[www.neorni.be](http://www.neorni.be)  
[www.neornilab.be](http://www.neornilab.be)



NEORNIVET

Veterinary practice



NEORNILAB

Laboratory

Gender • Performance genes  
Kinship • Autopsy  
Disease testing • Bacteriology  
Species determination •  
Faecal analysis • Colour  
mutations

NEORNI



PHARMA

BY VETS FOR BIRDS

Product range



Follow us  
on Facebook



Neorni (all services)

Beverlosesteenweg 129  
3583 Paal, Belgium  
T +32 (0)11 96 91 02

---



# BREEDING & IMMUNITY

# Natural breeding boost

## Basic breeding

The basic breeding schedule is the essence of successful breeding. Several field trials have shown that the birds were better prepared for breeding, which was reflected in more nests and more young per nest.



### Immuno-Plus

2 ml/l or 4 ml/kg

**Prior to breeding:**  
daily for a period of 6 weeks until 1st clutch

Breeding urge +++

Polypeptides

Stimulation of egg cell & sperm cell production



### Immuno Booster NEW

20 g/kg egg food/feed

**During breeding:**  
daily from hatching until weaning

Breeding urge +

Polypeptides & Premix breeding and rearing

Maintenance of egg cell & sperm cell production



### Need advice?

Your Neorni veterinarian will be happy to help



### Vit E + Se

2 ml/l or 4 ml/kg

Daily for 1-2w from breeding until the 1st clutch

Fertility +++

Linseed oil, arginine, spirulina, carnitine, zinc, etc

Extra high dose of vit E + Se

6

## Extra breeding (fertilisation)

Fertilisation problems occurring upon application of the basic schedules often have a nutritional or infectious cause.

►► nutritionally speaking, a **Vit E + Se** supplement could certainly offer added value.

►► In addition, certain germs with an important influence on fertilisation have been isolated in recent years. Faeces culture before breeding can be carried out easily and quickly at NeorniVet veterinary practice(± 24h).

►► In specific situations/species, extra added value can be achieved with the use of specific supplements such as **Vita-Plus, Ready, Hepa-Plus ...**

# IMMUNO BOOSTER NEW



## All-in premix for young birds



### Indication

- Highly concentrated premix
- Supports breeding results and maintains the condition of your birds
- Supply of extra energy
- Supports the immune system
- Helps build up intestinal flora
- Good development of the skeletal system
- Ideal for animals that are required to deliver top performances

### Administration 20 g/kg

- 1 large scoop (20 g)/kg egg food/feed or 2 small scoops (2 g)/100 g egg food/feed
- During breeding season: add daily to egg food/feed
- Extra energy & accommodating deficits: 1-2 days per week throughout the year

*Corn gluten, potato protein, potato protein fermentate, yeast and parts of it, chicory root fibre, spirulina, yeast, trimagnesium citrate, monoglycerides, diglycerides and triglycerides of fatty acids (c4), tricalcium citrate, L-carnitine, betaine, vitamin A, vitamin D3, vitamin E, vitamin C vitamin B1, vitamin B2, calcium D-pantothenate, vitamin B6, vitamin B12, vitamin K, niacinamide, folic acid, choline, clinoptilolite, copper, iron, zinc, manganese, potassium iodide, selenium, cobalt, lysine, methionine, glycine, echinacea purpurea, curcuma, lactobacillus acidophilus*





## Breeding, show/competition, extra energy



100 ml €25  
250 ml €55  
1 l €179

8

### Indication

- Supports breeding results and maintains the condition of your birds
- Supply of extra energy
- Good appetite and feather patterns
- To be used before/after intense exertion/competition
- Supports the development & growth of young animals

### Administration 2 ml/l drinking water

- Breeding: daily from 6 weeks prior to breeding. Stop administration from first clutch. After hatching, administer again daily to parents (and chicks) until fledging.
- Show/competition: starting 5 days before show/competition, daily 2 ml/l drinking water or 2 ml/kg body weight directly into the bill
- Supply of extra energy: 2 ml/l drinking water for a period of 14 days or 2 ml/kg body weight directly into the bill until recuperated



## Immunity, recovery, energy



80 capsules

€24

### Indication

- Stimulates recovery after illness/exertion
- Supply of extra energy
- Supports the immune system
- Ideal for top performance

### Administration

- Recovery/immunity boost: 1 capsule/0.5-1 kg orally for 5-10 days
- Extra energy: 1 capsule/0.5-1 kg orally
- Pigeon: day of return home, 1 capsule/pigeon
- To support breeding: for a period of 14 days before mating until laying 1 capsule/0.5-1 kg
- *Immuno Recup Caps* can be administered directly or with the food. When administered via food, the full dose should be consumed.

*Maize gluten, potato protein, potato protein fermentative, yeast and parts of it, chicory root fibre, spirulina, yeast, trimagnesium citrate, monoglycerides, diglycerides and triglycerides of fatty acids (c4), tricalcium citrate, L-carnitine, Vit A, Vit D3, Vit E, Vit C, Vit B1, Vit B2, calcium D-pantothenate, Vit B6, Vit B12, Vit K, niacinamide, folic acid, choline, clinoptilolite, copper, iron, zinc, manganese, potassium iodide, selenium, cobalt, lysine, methionine, glycine, echinaceae purpureae, curcuma, lactobacillus acidophilus*



## Start-up supplement for chicks, black spot



200 g €25

10

### Indication

- Smooth start for your young chicks in the first 6 weeks of life
- Natural suppression of bacterial and necrotic enteritis/diarrhoea
- Rapid growth and larger birds due to smooth start and natural suppression of black spot

### Administration 5 g/kg egg food/feed

- Daily throughout the first 6 weeks of life



# VIT E + SE



## Extra vitamin E + selenium



100 ml €17

250 ml €35

+ spirulina  
+ zinc  
+ linseed oil

### Indication

- For animals with a deficiency of vitamin E selenium
- Recommended for nervous symptoms

### Administration

- 2 ml/l drinking water or 1 ml/kg body weight directly into the bill
- Recommended for nervous symptoms: 7-10 days




#### Fertility +++

2 ml/l or 4 ml/kg feed daily for 1-2 weeks, prior to breeding until first clutch

# Fertility therapy

Individual birds (cocks, hens, psittacidae, etc)





# LIVER, MUSCLES & FITNESS



## Liver support



100 ml €20  
250 ml €36  
1 l €96

14

### Indication

- Supports liver function
- Relieves stress on the liver

### Administration

- Curative: 4 ml/l drinking water for a period of 5 days
- Prevention: 2 ml/l drinking water, 2 to 3 days per week



Can be administered daily to support the liver during administration of medication or dye



## Form, condition & improved blood flow



100 ml €22

250 ml €48

1 l €144

*Additional  
iron, iodine,  
L-carnitine &  
vit. B12*

### Indication

- Condition booster: improves flight performance
- Stimulates fat metabolism and endurance
- Supports blood circulation

### Administration 5 ml/l drinking water

- **Condition boost:**
  - For exhibition/flying season: for 2 weeks
  - For important flight: for 2-5 days
- **Fat birds:** for 3-5 days a week until the result is achieved
- **Anaemia:** 7-10 days
- Adjusted dosage songbirds <250 g body weight (canary, siskin, etc.): 1-2 ml/l

*Glycerol, L-carnitine, beetroot juice, potassium iodide, iron glycinate, vitamin B12, magnesium citrate sodium chloride, magnesium chloride, calcium chloride, glucose, vitamin E, selenium, zinc, copper, manganese, vitamin A, vitamin B1, vitamin B2, vitamin B5, vitamin B6, vitamin PP (niacinamide), vitamin D3, vitamin C, blend of natural aromatic components*

# Dosing card

## NEORNI

Beverloosesteenweg 129, Paal  
••• Tel. +32 (0)11 96 91 02 •••



Veterinary practice



Laboratory



Product range



## CAPSULES

- Alternative administration of capsules via drinking water (ml) or feed (g)
- For all cases where inserting is not possible

Capsule	/100 ml	/50 g
SBS Caps	1	1
Asper Dox Caps	1	1
Endo-Ecto Plus Caps 0.5 kg	-	1
Orni Caps 1 kg	-	1
Sulfaclozine Caps	1	1-2

Capsule	/300 ml	/150 g
Endo-Ecto Plus Caps 1 kg	-	1
Tricho Cox Caps	1	1



**NeorniPharma**  
powders & capsules

16



## POWDERS

- Easy administration of powders via drinking water (l) or feed (g/kg), without using a scale
- Level off\* the spoon after scooping

'PURE' Powder	
Amoxicillin 80%	1 / l
Colistin 480 IU/mg	2 / l
Doxycycline 15%	2 / l
Doxycycline 50%	1 / 2 l
Fungizone powder/Ampho B	1 / l
Lincospectin 66%	1 / 1.5 l
Spiramycin 50%	1 / 1.5 l
Tiamulin 45%	1 / 2 l
Trim-Sulfa 90%	1 / 1.5 l
Tylosin 100%	2 / l

\*Spoon included with your purchase



Powder	
Airfree	4 / l
Bronchifree	3 / l
CAS (loft disinfection)	3 / l
Coli-Mix	3 / l
Gould Mix	3 / l
Immuno Booster NEW	2 / 100 g
Megadysbac	3 / l
Mucolyt	3 / l
Neocox	1 / 150 g
Probi-Plus	2 / l
Red Race	2.5 / 2 l
Sulfaclozine	2 / l
Super Start-Up	4 / kg
Tricho Geel	3 / 2 l (<0.5 kg: 1 / l)
Vita Energy	5 / kg or 2 / l
YPD Mix	3 / l



### Visit us

NeorniPharma  
Beverloosesteenweg 129  
3583 Paal, Belgium

### Contact us

T +32 (0)11 96 91 02  
E info@neornipharma.be  
W www.neornipharma.be

Download our catalogues




Follow us on Facebook



**NeorniPharma, formulations tailored to your birds**

The dosages on this card offer an alternative administration compared to the product label, tailored to your bird or situation. Contact us if in doubt.





# MEGABACTERIA & DYSBACTERIOSIS

# Dysbacteriosis

crop

proventriculus

intestines



**Dysbac Liquid**

prevention of  
dysbacteriosis

**7-in-1**

Intestinal health  
on 7 fronts

**Dysbac Liquid** is an improved & extended version of **CU-For Liquid** with more focus on **intestinal health** and **general immunity**

18

2 ml/l
+
+++
+++
+++
+++
+++
<b>CROP, PROVENTRICULUS &amp; INTESTINAL HEALTH, PREVENTION OF DYSBACTERIOSIS</b>

- **Short-chain fatty acids:** formic acid, propionic acid, citric acid
- **Medium-chain fatty acids:** butyric acid, monoglycerides, diglycerides and triglycerides of fatty acids (C8-C10)
- **Aromatic substances:** oregano, echinaceae purpureae, ginger, cinnamon, black pepper, garlic, and boswellia
- **Vitamin C**
- **Chelated copper and zinc**
- **Extra fibre**
- **Fish oil**

DOSAGE
ACIDIFICATION OF DRINKING WATER (pH)
CROP HEALTH
POVENTRICULUS HEALTH
INTESTINAL HEALTH
GENERAL IMMUNITY
PURPOSE

1-2 ml/l
+++
+++
++
-
-
<b>WATER ACIDIFICATION, WATER QUALITY</b>

- **Short-chain fatty acids:** ammonium formate, acetic acid, lactic acid, formic acid, propionic acid
- **Copper and zinc sulphates**



**Need advice?**

Your Neorni veterinarian will be happy to help





## Crop, proventriculus and intestinal health



### Indication

- Forms a protective film over the intestinal mucosa
- Increases the animal's immunity
- Helps to maintain the natural balance of intestinal flora
- To be used in cases of watery droppings

### Administration 2 ml/l drinking water

- Prevention: 1 to 2 days per week
- In case of digestion problems: for a period of 7-14 days



### Prevention of megabacteriosis & dysbacteriosis

- 1-2 days a week 2 ml/l drinking water
- May be combined with ACS

# SilverSpray

**New** disinfectant spray

Bird-friendly  
Broad-spectrum  
Ready-made  
Multi-purpose



---



# NATURAL OILS



## Flight oil, source of energy from lipids



22

### Indication

- Source of energy from lipids
- Feather quality, optimal feathering
- Extra vitamins A, D, and E
- Hoge source of omega 3-6-9 fatty acids

### Administration 5-10 ml/kg feed

- During moulting: 2 days per week
- Before and during breeding: 2-3 days per week
- During racing season: 1-2 days per week
- *Feather Energy* can be combined with *Vita Energy* or *Immuno Booster NEW*



Can be used to improve the absorption of *Immuno Booster NEW* and *Vita Energy* with the feed



## Unobstructed respiratory tract



30 ml €17

### Indication

- Hardened snot and black nares

### Administration 1-2 drops per naris

- All birds: for respiratory problems: 2-3 times/day for 7 days



## Pecking, restlessness



NEW

100 ml €28



24

### Indication

- Pecking in birds and poultry

### Administration

- 1 ml/l drinking water for 2-4 weeks



#### Feather picking/pecking

Administer simultaneously with *Immuno-Plus* for the best results



## Respiratory tract, heat stress



100 ml €19

250 ml €35

formerly  
**Herbi+E**

### Indication

- Natural solution to support respiratory problems
- Suppression of pain/inflammation in case of respiratory problems
- Stuck slime
- Heat stress

**Administration** 3 ml/l drinking water or 0.5 ml/kg body weight directly into the bill

- Preventive: when purchasing new birds: for 10-14 days
- Curative: in mild to moderate respiratory problems, for 10-14 days
- Heat stress: 3-5 ml/l



**Additional support for racing pigeons in hot weather** 3 ml/l of drinking water, in combination with *Aminovital* on the basketing day

*Vitamin A, vitamin C, curcuma, sage, elderberry, thyme, oregano, mint, eucalyptus, echinacea, grindelia, clove, nettle, rosemary*

NEW

# neofalco

products for falcons



more info: [info@neornipharma.be](mailto:info@neornipharma.be)





# PARASITES



## Red mite, northern fowl mite



50 ml €35

28

### Indication

- Spray for surroundings (coop) and birds to combat haematophagous mites
- Not harmful to birds: spray may be applied directly to birds; coop may be treated in the presence of the birds

### Administration 8 ml/l/40 m<sup>2</sup>

- Spray: 8 ml/l water for 40 m<sup>2</sup> coop
- Repeat after 1 week



## Ectoparasites



250 ml €22

1 | €69

### Indication

- Ectoparasites

### Administration

- Red mite/feather lice/northern fowl mite: Spray at the level of the neck, tailbone and cloaca, and under the wings. Once, repeat after 7 days.
- Scaly leg mites: Spray the area to be stabilised daily for 14 days.
- Feather mite: Spray at the level of the neck, tailbone and cloaca, and under the wings. Once, repeat after 10 days and after 20 days.
- Preparation for breeding: Preventively spray nests/ breeding boxes/coops/perches with *Exmite Spray*. May be repeated weekly in case of high load.



#### Heavy feather mite infections

Day 0, 10 and 20: serve simultaneously *V-Exmite Drops* and/or *Endo-Ecto Caps* for best results

## Notes

[illegible]

---



# PROBIOTICS & DISINFECTION

# NEORNI STABILIZER



## Probiotic ambient spray



1 | €20

32

### Indication

- Reduction of pathogenic load in your birds' surroundings
- Forms a stable and healthy environment by occupying it with safe, probiotic bacteria

### Administration

- To be used diluted: 5-10% (5-10 ml/100 ml) in lukewarm water
- Mist the entire area to be stabilised
- First week, spray the coop daily (may be done in the presence of birds, including nestlings).
- Subsequently, spray twice a week to maintain probiotic protection
- The product diluted with water has a maximum shelf life of 5 days.



Can be nebulised in the presence of the birds/pigeons, especially during important or stressful periods when the bacterial presence in the environment rises



## Nasal cleaner



NEW

50 ml €10

### Indication

- Cleans the nares
- Loosening and removing trapped mucus and dust in the upper respiratory tract

### Administration 1-2 drops/naris

- The snot is shaken off by the pigeons after administration. If not, manually remove the snot after 5 minutes.
- In case of problems: daily or every other day
- Breeding pigeons: 14 days before mating
- Flying pigeons: 14 days before the first flight
- Juvenile pigeons: 14 days before first training and first flight
- Important flights: 1 week before basketing



## Probiotics and prebiotics



34



### Indication

- Increases the beneficial flora in the crop and the intestines
- Ensures improved digestion and supports the initiation of the crop and intestinal flora in chicks
- After a course of antibiotics

### Administration 3 g/l drinking water or 6g/kg feed

- Preventive: for 2 days/week
- Wet droppings or after antibiotic treatment: for 5-7 days
- Refresh daily
- May be combined with other supplements (vitamins)



Change drinking water twice daily or administer through the feed





## Disinfectant spray



### Indication

- Broad-spectrum disinfectant
- Bactericide - Fungicide - Virucide

### Administration

- Ready-to-use disinfectant spray
- Suitable for pens, cages, equipment and eggs



Safe to apply in the presence of your birds



## Probiotic spray for skin infections



200 ml €17

36

### Indication

- Suitable for birds with sensitive skin
- To be used externally for care of the skin
- Reduces the formation of scaly skin


### Administration

- Mist the area to be stabilised



#### **Powder comb/flavus/white flakes**

Spray the comb or infected area daily. Not harmful to eyes or respiratory tract.



# TRICHOMONIASIS, CANDIDA ASPERGILLOSIS & DRINKING WATER DISINFECTION



## Acidification of drinking water



250 ml €15

38

### Indication

- Acidification of drinking water, lowering pH

### Administration 1-2 ml/l drinking water

- Prevention: 1 to 2 days per week



Exceptionally concentrated formula, very economical to use



## Drinking water disinfection



NEW

1 | €32

### Indication

- Water disinfection
- Prevention of harmful bacteria, viruses, yeasts and moulds
- Immunity support

### Administration

- 2 ml/l drinking water



Longer and better action due to silver addition



## Throat elixir



50 ml €17

250 ml €59

1 l €165

40

### Indication

- Preventive approach to trichomoniasis
- Supports throat and crop health
- Positive effect on the upper respiratory tract
- Plays a role in resistance to disease

### Administration

- The day after returning home:
  - Recommended administration: 10 ml/kg feed
- **Always make fresh before each feeding.**
- **Most effective up to 2 hours after mixing.**
  - 5 ml/l drinking water
  - 1 drop per naris
- Prevention: 1 day per week.
- At high pressure: 2 days per week
- Caution: stains clothing and hands



Proven efficacy in resistant trichomoniasis cases:  
for 5 days 10-15 ml/kg feed

---



# VITAMINS, AMINO ACIDS, ELECTROLYTES, MINERALS & TRACE MINERALS





## Amino acids, electrolytes & vitamins



100 ml €19

250 ml €35

1 l €65

Extra vitamin E,  
selenium & BCAA  
amino acids

42

### Indication

- All essential amino acids crucial for proper moulting/feather development/pigmentation
- Stimulates recovery/repair
- Muscle building and recovery

### Administration 2 ml/l drinking water

- Racing pigeons: 5 ml/l during the racing season
- Throughout the entire year: 1 to 2 days per week
- Throughout the entire moulting period: 3 days/ week. May be combined with *Condi-Plus*.



**Additional support for racing pigeons in warm weather** 5 ml/l drinking water, in combination with *Respicor* on the basketing day

*Dextrose, glycerol, sodium chloride, prebiotic fibre, trimagnesium citrate, beetroot juice, potassium chloride, sucrose, calcium chloride, tricalcium citrate, lysine, methionine, threonine, tryptophan, histidine, carnitine, betaine, choline, vitamin B1, vitamin B2, niacinamide, calcium D-pantothenate, vitamin B6, vitamin B12, folic acid, biotin, vitamin C, vitamin E, selenium, copper, zinc, manganese, mix of natural aromatic components, L-isoleucine, L-valine, L-leucine*





## Moulting supplement



100 ml €15  
250 ml €26  
1 l €83

### Indication

- Supports bird health during moulting
- Supports the metabolism
- Supply of extra energy
- Increases appetite

### Administration 2 ml/l drinking water

- Throughout the entire moulting period: 3 days/ week. May be combined with *Aminovital*.



## Conditioning powder – Vitamins, minerals and amino acids



44

### Indication

- Condition booster & extra energy
- Extra vitamins, minerals and amino acids
- Increases endurance
- Breeding stimulation

### Administration

- Recommended administration: 10 g/kg feed
- Administration via drinking water: 4 g/l



- General: Throughout the year, 1-2 days/week.  
In case of poor/impaired condition: for a period of 14 days.
- Breeding stimulation: 2-4 weeks prior to breeding, twice a week to daily
- Pigeon: Medium distance/long distance, 2-5 days before basketing

*Dextrose, maltodextrin, soya protein concentrate, calcium carbonate, yeasts and parts thereof, trimagnesium citrate, monosodium phosphate, sodium bicarbonate, sodium chloride, potassium chloride, trisodium phosphate, tricalcium citrate, vitamin A, vitamin D3, vitamin E, vitamin K, vitamin B1, vitamin B2, vitamin B3, vitamin B5, vitamin B6, vitamin B11, vitamin B12, biotin, vitamin C, choline chloride, iron, copper, zinc, manganese, potassium iodide, selenium, lysine, methionine, tryptophan, threonine, mixture of natural aromatic components*



## Liquid source of vitamin D, calcium, phosphorus and other minerals



100 ml €17

250 ml €32

1 l €88

Extra calcium  
& phosphorus

### Indication

- For birds with a deficiency in Vitamin D3, calcium, phosphorus and other minerals
- Supports bone formation and development in young animals
- Supports egg shell formation
- Poor egg shell quality/shell-less eggs

### Administration 2 ml/l drinking water

- Prevention: 2 ml/l
  - 2 weeks prior to breeding until the eggs are laid: 2-3 days per week
  - Throughout the first 6 weeks of life: 2-3 days per week
- Curative: 5 ml/l for a period of 7 days



Change drinking water twice daily to avoid sedimentation, or administer via the feed (preventive: 4 ml/kg egg food/feed - curative: 10 ml/kg egg food/feed)

*Trisodium phosphate, trimagnesium citrate, calcium chloride, magnesium chloride, tricalcium citrate, vitamin D3, cholecalciferol, zinc, manganese, copper, iron*



## Water-soluble multivitamins



46

### Indication

- Supplement containing all essential vitamins
- Overall daily health
- Prevention of chronic vitamin A deficiency

### Administration 2 ml/l drinking water

- Weekly: for 1 to 2 days per week
- After illness/recuperation: for a period of 3 to 5 consecutive days



For parrots and parakeets, we strongly recommend 1-2 days per week *Administer Vita-Plus* via the drinking water to prevent vitamin A deficiency



High concentration of vitamin K helps prevent or stop bleeding

*Vitamin A, vitamin D3, vitamin E, vitamin K3, vitamin C, vitamin B1, vitamin B2, vitamin B6, vitamin B12, vitamin PP, biotin, folic acid, sorbitol*

---



# PURE PURE

# BREWER'S YEAST



## Indication

- Amino acids, minerals and trace elements
- Vitamin B

**Administration** 15-30 g/kg feed

1 kg €18

# DEXTROSE



## Indication

- Additional energy source
- Flavouring medicated water

**Administration** 15-30 g/l drinking water

1 kg €18

# ELECTROLYTE



## Indication

- Hydration in case of dehydration
- Stabilisation of water and electrolyte balance

**Administration** 15-30 g/l drinking water

500 g €18

48

# YELLOW



## Indication

- Colour

**Administration** 1-2 g/kg feed

100 g €15

500 g €60

# HIGHLY DIGESTIBLE PROTEIN



## Indication

- Muscle growth
- Additional amino acids/proteins

**Administration** 5-15 g/kg feed

400 g €25

# RED



## Indication

- Colour

**Administration** max. 1 g/kg feed

100 g €28


500 g €85



# SCHEDULES FOR SONGBIRDS

# Annual Euro & Canary schedule

## Timeline



### Preparation for breeding

- Breeding boost
- Liver support
- Black spot prevention

### Hatching to weaning

- Optimal start & mushy chick disease prevention
- Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi

### Weaned young

- Prevention coccidiosis/atoxoplasmosis, megabacteriosis and dysbacteriosis
- Limiting pressure with immunity build-up







# Annual schedule for Euro & Canary



*The annual schedule is a general guide for preparation, breeding, hatching, weaning and moulting. Talk to your veterinarian about the specific requirements for your birds and devise a tailored schedule together.*

## Preparation from 6-8 weeks prior to breeding

1. A healthy environment is the best start! (see CAS or SilverSpray)
2. Prevention of atoxoplasmosis/coccidiosis + megabacteriosis/dysbacteriosis
  - **Veterinarian:** bird check-up + faecal analysis 6-8 weeks prior to breeding
  - Only use courses of treatment in case of problems:
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 3 days Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
3. For a period of 6 weeks prior to breeding
  - **Immuno-plus (2 ml/l) daily**
    - Breeding boost; birds will be in top condition
  - 1-2 days per week **Hepa-Plus (4 ml/l)**
    - Liver support
  - As required: Vita-Plus, Vita Energy, Ready, Vit E+Se, Vita-Min-DCP
    - These are suitable for use with Immuno-Plus
  - Extra: black spot prevention
    - For 7 days **Gould Mix (5 g/l)** during breeding preparation
4. Prevention of haematophagous mites
  - **Exzolt:** 1 ml/l for 1 day, repeat on days 7 and 14
    - Does not affect fertility & nestlings/young
    - Can be repeated every 3 months
  - Extra prevention of haematophagous mites
    - Before occupation, treat coop/loft with:
      - **Derma-Plus/Exmite Spray**
      - Can be repeated while the birds/nestlings/young are present

51

## Breeding: hatching to weaning

1. **Immuno Booster NEW (2 g/100 g or 20 g/kg egg food/feed)**
  - All-in premix for young birds
2. First 6 weeks
  - Option 1: **Super Start-Up (5 g/kg egg food/feed)**
    -  Natural suppression and prevention of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
  - Option 2: **Super Start-Up Blue (20 g/kg egg food/feed)**
    -  Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
3. Start from 2 days before hatching until 5 days after hatching **Coli-Mix (5 g/l)**
  - Prevention of mushy chick disease (Colibacillosis)
  - Alternative to Coli-Mix: **Gould Mix (5 g/l)**



## Breeding: weaned young

1. A healthy environment is the best start!

- **CAS or SilverSpray**
  - Environment free of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and moulds
- **Neorni Stabilizer:** Probiotic ambient spray
  - Prevention by reducing bacterial presence in the environment

2. Preventive cure

- 3rd week after weaning
  - **Increased susceptibility to atoxoplasmosis/megabacteriosis**
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 3 days Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)

3. Then throughout the year (from weaning to breeding period)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis

## Moulting period

1. Throughout moulting **Condi-Plus + Aminovital:**

- Condi-Plus (2 ml/l): 3 days/week
- Aminovital (2 ml/l): 3 days/week
- May be administered simultaneously

2. Throughout the year (from weaning to breeding)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis

At NeorniPharma, we always strive for the best long-term solution. We are constantly looking for **alternatives to anti-microbials** to limit the risk of resistance. The goal is to give our birds **as few preventive treatments with antimicrobial agents as possible**. Clinical research and any additional tests are crucial for this. After the diagnosis, an appropriate treatment can be administered. **Antibiotics are not harmless**. Consult your vet for advice about their correct and responsible use. **Autopsy:** don't simply throw away a bird that has died! The cadaver can provide a lot of information to ensure appropriate treatment can be sought. It is best to keep the bird cool (in a fridge, for example) and carry out an investigation within 24 hours (max. 72 h).



# Preparation for breeding



**From 6 weeks  
prior to breeding  
until the  
1st clutch**



## **Need advice?**

Your Neorini  
veterinarian will be  
happy to help

### **Immuno-Plus**

2 ml/l

6 weeks prior to mating  
until the 1st clutch

Daily

Stimulation of sperm cell &  
egg cell production

### **Hepa-Plus**

4 ml/l

6 weeks prior to mating  
until the 1st clutch

1-2 days/week

Supports the liver

53



**Extra  
support**

### **Vit E + Se**

2 ml/l

2 weeks prior to mating  
until the 1st clutch

Daily

Extra-high doses of Vit E +  
Se, linseed oil, arginine, spi-  
rulina, carnitine, zinc, etc.

### **Vita-Min-DCP**

2 ml/l

2 weeks prior to mating  
until the 1st clutch

2 days/week

Extra calcium, phosphorus,  
vit D3, and trace minerals

### **Ready**

3-5 ml/l

6 weeks prior to mating  
until the 1st clutch

1-2 days/week

Converts fat into usable  
energy for the bird

### **Vita-Plus/Vita Energy**

2 ml/l or 10 g/kg

A: Throughout the entire year;  
B: 2 weeks before mating until  
the 1st clutch

A: 1-2 days/week  
B: Daily

Extra vitamins

**FERTILISATION**

**EGGSHELL**

**CONDITION**

**VITAMINS**

▶▶▶▶ All products can be used in combination





# Breeding: hatching to weaning



54

## Coli-Mix

5 g/l or 10 g/kg egg food/feed

2 days before to 5 days after hatching

Does not affect fertility

Broad-spectrum mix

**Prevention of mushy chick disease & Bengalese finch disease**

## Immuno Booster NEW

20 g/kg egg food/feed

Daily from hatching until weaning

Positive effect on fertility

All-in premix

**Maintenance of egg cell & sperm production**

## Super Start-Up

5 g/kg egg food/feed

**Natural suppression and prevention of coccidiosis/ atoxoplasmosis, bacteria, viruses, yeasts and fungi**

**OR**

## Super Start-Up Blue

20 g/kg egg food/feed

**Suppression of coccidiosis/ atoxoplasmosis, bacteria, viruses, yeasts and fungi**

**Immuno Booster NEW** contains all essential components for **rearing young birds**. It is a unique **comprehensive** premix to mix under egg food/feed, providing all necessary nutrients in **exceptionally high concentrations**.

**Immuno Booster NEW Polypeptides** Keep the birds in form during breeding. **Saccharomyces cerevisiae** Source of protein, amino acids, and nucleotides. **Chicory root fibre** Prebiotic fibre: balanced digestion. **Spirulina** Source of protein, fats, carbohydrates, vitamins, and minerals. **Electrolytes** Fluid balance. **Fatty acids** Energy, balanced digestion. **All water-soluble and fat-soluble vitamins** General immunity, antioxidants, growth, energy release. **All amino acids** Building muscle and feather cover. **L-carnitine** Supports fat digestion and muscle building. **Choline** Liver support. **Minerals** (cobalt, iron, copper, zinc, manganese, iodine, selenium) General immunity, antioxidants, growth, energy release. **Clinoptoplite** General health and intestinal health. **Probiotics** Intestinal flora stabiliser. **Turmeric and echinacea** General immunity and health.



# Breeding: weaned young

COCCIDIOSIS & ATOXOPLASMOSIS	MEGABACTERIA & DYSBACTERIOSIS
<b>ACS</b>	<b>Dysbac Liquid</b>
0.5 ml/l	2 ml/l
From weaning until next breeding time	From weaning until next breeding time
3-5 days/month If required: twice monthly	1-2 days/week
<b>COCCIDIOSIS &amp; ATOXOPLASMOSIS</b>	<b>MEGABACTERIOSIS, DYSBACTERIOSIS &amp; FLAGELLATES</b>



## Extra support throughout the year

SUPPORTS THE LIVER	GENERAL SUPPORT
<b>Hepa-Plus</b>	<b>Vita-Plus / Vita Energy</b>
2 ml/l	2 ml/l or 10 g/kg
Throughout the year	Throughout the year
1-2 days/week When adding dye: 2 ml/l daily	1-2 days/week
<b>LIVER SUPPORT</b>	<b>EXTRA VITAMINS</b>



# Annual Exotics schedule





*The annual schedule is a general guide for preparation, breeding, hatching, weaning and moulting. Talk to your veterinarian about the specific requirements for your birds and devise a tailored schedule together.*

## Preparation from 6-8 weeks prior to breeding

1. A healthy environment is the best start! (see CAS or SilverSpray)
2. Prevention of flagellates (cochlosomosis) + campylobacteriosis
  - **Veterinarian:** bird check-up + faecal analysis 6-8 weeks prior to breeding
  - Only use courses of treatment in case of problems:
    - For 7 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)
3. For a period of 6 weeks prior to breeding
  - **Immuno-Plus (2 ml/l) daily**
    - Breeding boost; birds will be in top condition
  - 1-2 days per week **Hepa-Plus (4 ml/l)**
    - Liver support
  - As required: Vita-Plus, Vita Energy, Ready, Vit E+Se, Vita-Min-DCP
    - These are suitable for use with Immuno-Plus
  - Prevention of cochlosomosis + campylobacteriosis
    - For 7 days **Gould Mix (5 g/l)** during breeding preparation

4. Prevention of ectoparasites
  - Haematophagous mites
    - **Exzolt:** 1 ml/l for 1 day, repeat on days 7 and 14
      - Does not affect fertility & nestlings/young
      - Can be repeated every 3 months
    - Extra prevention of haematophagous mites
      - The day prior to occupation, treat coop/loft with:
        - **Derma-Plus/Exmite Spray**
        - Can be repeated while the birds/nestlings/young are present
  - Tracheal mites
    - **Vermi-Complete:** 2.5-5 ml/l for 1 day, repeat after 3 weeks

## Breeding: hatching to weaning

1. **Immuno Booster NEW (2 g/100 g or 20 g/kg egg food/feed)**
  - All-in premix for young birds
2. First 6 weeks
  - Option 1: **Super Start-Up (5 g/kg egg food/feed)**
    -  Natural suppression and prevention of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
  - Option 2: **Super Start-Up Blue (20 g/kg egg food/feed)**
    -  Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
3. Start from 2 days before hatching until 5 days after hatching **Coli-Mix (5 g/l)**
  - Prevention of mushy chick disease (Colibacillosis)
  - Alternative to Coli-Mix: **Gould Mix (5 g/l)**



#### 4. For problems in the first 6-8 weeks of life

- **Gould Mix (5 g/l)**
  - For the treatment of cochlosomosis (Bengalese finch disease), yeasts and bacterial diarrhoea

### Breeding: weaned young

#### 1. A healthy environment is the best start!

- **CAS or SilverSpray**
  - Environment free of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and moulds
- **Neorni Stabilizer:** Probiotic ambient spray
  - Prevention by reducing bacterial presence in the environment

#### 2. Preventive cure

- 3rd week after weaning
  - **Increased susceptibility to coccidiosis/cochlosomosis/campylobacter**
    - 5 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)
    - 3 days Hepa-Plus (4 ml/l)
    - 5 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)

#### 3. Then throughout the year (from weaning to breeding period)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis
  - Prevention of flagellates (cochlosomosis)

57

### Moulting period

#### 1. Throughout moulting **Condi-Plus + Aminovital:**

- Condi-Plus (2 ml/l): 3 days/week
- Aminovital (2 ml/l): 3 days/week
- May be administered simultaneously

#### 2. Throughout the year (from weaning to breeding)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis
  - Prevention of flagellates (cochlosomosis)

At NeorniPharma, we always strive for the best long-term solution. We are constantly looking for **alternatives to anti-microbials** to limit the risk of resistance. The goal is to give our birds **as few preventive treatments with antimicrobial agents as possible**. Clinical research and any additional tests are crucial for this. After the diagnosis, an appropriate treatment can be administered. **Antibiotics are not harmless**. Consult your vet for advice about their correct and responsible use. **Autopsy:** don't simply throw away a bird that has died! The cadaver can provide a lot of information to ensure appropriate treatment can be sought. It is best to keep the bird cool (in a fridge, for example) and carry out an investigation within 24 hours (max. 72h).

# Trichoryza Drops

The best way to tackle  
trichomoniasis  
**without losing form**

**FLYING SEASON**  
10 ml/  
kg feed  
1 day/week, day after returning home

**NEW**  
1 litre



**Proven effectiveness  
against resistant  
trichomoniasis!**





# SCHEDULES FOR PIGEONS



# Breeding



## 2 weeks prior to mating until laying

Trichoryza Drops	Vita-Min-DCP	Immuno-Plus
10 ml/kg feed	2 ml/l	2 ml/l
During brooding	2 weeks prior to mating until the 1st clutch	2 weeks prior to mating until the 1st clutch
For a period of 3 consecutive days	2 days/week	Daily
Proven effectiveness in resistant trichomoniasis	Extra calcium, phosphorus, vit D3, and trace minerals	Stimulation of sperm cell & egg cell production

## During brooding

60

Tricho Geel
3 g/l or 6 g/kg feed
During brooding
For a period of 5 consecutive days
Reduce pressure of trichomoniasis + avoid hexamitiasis in youngsters



## From hatching to weaning

Immuno Booster NEW + Feather Energy	Trichoryza Drops	Aminovital
20 g/kg feed + 5 ml/kg feed	10 ml/kg feed	2-5 ml/l
From hatching to weaning	Full breeding period	From hatching to weaning
2-3 days/week	For 1 day every 2 weeks	3 days/week
All-in premix	Proven effectiveness in resistant trichomoniasis	Additional amino acids and vitamins for rapid growth



# Flight preparation

## OLD PIGEONS AND YEARLINGS



**Trichoryza Drops**



**Dysbac Liquid**



**Ready**



10 ml/kg



1 day/2 weeks



Proven efficacy in resistant trichomoniasis



Insufficient training: **Immuno-Plus (2 ml/l) + Ready (2 ml/l)** for a period of 14 days

2 ml/l

2 days/week

Crop, proventriculus & intestinal health, solid droppings

5 ml/l

Daily for 2 weeks before the start of the racing season

Form & fitness building

## YOUNG PIGEONS 'JUNIOR'

61



**Immuno Booster NEW**



**Trichoryza Drops**



**Dysbac Liquid**



20 g/kg



2-3 days/week for 6-8 weeks, afterwards 1 day/week throughout flying season



Growth & energy, prevents nutritional deficiencies



Insufficient training: **Immuno-Plus (2 ml/l) + Ready (2 ml/l)** for a period of 14 days

10 ml/kg

1 day/2 weeks

Proven efficacy in resistant trichomoniasis

2 ml/l

2 days/week

Crop, proventriculus & intestinal health, solid droppings



### Need advice?

Your Neorni veterinarian will be happy to help

►►► This schedule is an indication of how the NeorniPharma products can be used in certain circumstances. We try to develop a complete and appropriate schedule with as few products as possible. All these supplements can be used in combination, if required. In case of poor performance, we will look for the cause with you and decide upon any adjustments or further treatments. Attention: food supplements are best prepared the day before so they can be well absorbed, which will help the pigeons digest the food better.





# Flying season

## LOW SPEED



Trichoryza Drops



Dysbac Liquid



10 ml/kg



Day after return home



For 1 day



Proven efficacy in resistant trichomoniasis



Pinch naris, add drop, allow bird to inhale

2 ml/l

Start of the week

For 2 days

Crop, proventriculus & intestinal health, solid droppings

Can be administered daily

62

## HIGH SPEED - MEDIUM DISTANCE



Aminovital



Trichoryza Drops



Dysbac Liquid



Ready



5 ml/l

10 ml/kg

2 ml/l

5 ml/l



Return home

Day after return home

Start of the week

Before basketing



For 1 day  
Demanding flights: for a period of 2-3 days

For a period of 1-2 days

For 2 days

Great speed: 1 day  
Medium distance: 2-3 days



Fast recovery

Proven efficacy  
in resistant  
trichomoniasis

Crop, proventriculus &  
intestinal health, solid  
droppings

Form



In hot weather:  
basketing day

Pinch naris, add drop,  
allow bird to inhale

Can be administered  
daily

Before an important  
race: daily for a period of  
10 days in combination  
with Immuno-Plus



# LONG DISTANCE

## Recovery



**Immuno Recup Caps**



**Aminovital**



**Immuno Booster NEW**



1 capsule/pigeon

5 ml/l

20 g/kg



For a period of 1-2 days

For a period of 2-3 days

For 2 days



Fast recovery

Fast recovery

Quick recovery, replenishing nutritional deficiencies

## Maintenance



**Trichoryza Drops**



**Dysbac Liquid**



**Respicor**



10 ml/kg

2 ml/l

3 ml/l



For 2 days

For 2 days

For 2 days



Proven efficacy in resistant trichomoniasis

Crop, proventriculus & intestinal health, solid droppings

Respiratory tract, white noses

63

## Boost before basketing



**Feather Energy**

**Vita Energy**

**Immuno-Plus**

**Red Race**



5 ml/kg

10 g/kg feed

2 ml/l

5 g/2 l



For a period of 3-5 days

For a period of 2-5 days

For a period of 3-5 days

For a period of 3-5 days



High-fat energy

Stimulates fitness

Extra energy

Additional nutrient uptake



# Moulting Support



## Aminovital

2 ml/l

3 days/week

**AMINO ACIDS +  
VITAMINS**



## Immuno Booster NEW

20 g/kg

2 days/week

**ALL-IN PREMIX**



## Feather Energy

5 ml/kg

2 days/week

**FATTY ACIDS**



## Treating ectoparasites



### V-Exmite Drops

1-2 drops in the neck

Moulting period

**INDIVIDUAL  
TREATMENT**

### Vermi-Complete

1 ml/l bath water

Moulting period

**GROUP TREATMENT**



64

## Preparation for breeding

Once the pigeon season finishes at the end of September/beginning of October, we strongly recommend you **collect mixed faeces from the breeders for a period of 5 days and have it analysed**. This mixed faeces is checked for worms, coccidiosis, and yeasts/fungi, following which a culture is grown to check for paratyphoid/salmonella. The results can be used to draw up a vaccination plan.

	Paratyphoid/salmonella NEGATIVE	Paratyphoid/salmonella POSITIVE
<b>Treatment</b>	Not necessary	Required for a period of 10 days according to antibiogram
<b>Vaccination</b>	Strongly recommended	Required
<b>Time of vaccination</b>	4 weeks before mating	2 days after treatment



### Need advice?

Your Neorni veterinarian will be happy to help

**Discuss a plan of action tailored to your coop and/or problem with our veterinary practice NeorniVet (+32 (0)11 96 91 02)**

### Faecal analysis

- Collect faeces every day for 5 days and refrigerate it.
- Send the faeces, or take your sample to your veterinarian
- If required, the pigeons can be checked at the same time
- This is not harmful to moulting pigeons





# SCHEDULES FOR PSITTACIDAE



# Psittacidae

## PARROTS & PARAKEETS

### General health



To prevent  
chronic  
deficiencies

Throughout the year  
1-2 days/week  
**Vita-Plus or Vita-Energy**

### Breeding

#### PREVENTION

6-8 weeks  
prior to  
breeding

Control mixed faeces from 3-5 days (vet)

- If necessary, deworm with **Vermi-Complete**
  - Group: 5 ml/l drinking water
  - Individual: 1 ml/kg body weight in the mouth

Control ectoparasites

- Group: **Vermi-Complete** 1 ml/l bath water
- Individual:
  - <250 g: **Endo-Ecto Kanarie**
  - >250 g: **V-Exmite Drops**

#### STIMULATES

daily 4-6  
weeks before  
breeding

**Vita Energy + Immuno-Plus**  
With drinking water and/or  
feed



#### REARING

daily from  
hatching to  
autonomy

Natural brooding:

- **Immuno Booster NEW** 2 g/100 g egg food/feed +  
**Immuno-Plus** 0.2 ml/100 ml drinking water

Hand rearing:

- **Immuno Booster NEW** 2 g/100 g soft food once per  
day + **Immuno-Plus** 1 ml/kg body weight once per day





# Crop and digestive problems



**For a period of 7-14 days**  
1 ml/kg body weight  
**Hepa-Plus**



**For 7 days:**  
1 capsule/kg body weight  
**SBS Caps**



**Need advice?**  
Your NeorNI  
veterinarian will be  
happy to help

67

## Veterinarian check-up

- Testing for disease through blood - feathers - cloacal swab
- Crop swab

# Moulting - Feathers



Additional  
support for  
spring growth

**Moulting + feather loss**  
3 days/week  
**Aminovital + Condi-Plus**  
2 ml/l 2 ml/l

## Notes

[illegible]



# SCHEDULES FOR CHICKENS

# Vaccination schedule



## Chicks

### Support

As required:

- First 14 days of life **Immuno Booster NEW** (20 g/kg feed)
- General vitamins: 3 days/week **Vita-Plus** (2 ml/l)
- Intestinal health + prebiotics: 2 days/week **Dysbac Liquid** (2 ml/l)
- Extra growth + amino acids: 2 days/week **Aminovital** (2 ml/l) + **Immuno-Plus** (2 ml/l)

### Vaccinations

**GOLD STANDARD:** 3 vaccines (Marek, NCD and smallpox)

D1 (1st week of life)	Marek (2 strains) + IBD + NCD (injection)
D1 (1st week of life)	Infectious bronchitis (eye drop)
1st week of life	Coccidiosis (drinking water)



## Growing chickens (from 7 weeks) & full-grown chickens

70

### Support

- General vitamins: **Vita-Plus** (2 ml/l) Prevention: 2 days/week After treatment: for 7 days.
- Intestinal health + prebiotics: **Dysbac Liquid** (2 ml/l). Prevention: 2 days/week In the case of flatter faeces: for 7 days.
- Respiratory tract support: **Respikor** (3 ml/l). Prevention: 2 days/week In the case of respiratory problems: for 7-14 days.

### Vaccinations

May-June (annual!)	Smallpox (+ ILT)
May-June (annual!)	NCD (injection/drinking water)
August-September	NCD (injection oil) <b>Mandatory for exhibition &amp; sale</b>
August-September	Infectious bronchitis (drinking water)

## Advice all ages

- **GOLD STANDARD:** 3 vaccines (Marek, NCD and smallpox). Other vaccines are relevant to specific problem areas, and not routinely recommended.





# Advice & tips

## General health



**Throughout the year**  
1 day/week  
**Vita-Plus (2 ml/l)**

**During moulting**  
3 days/week  
**Aminovital and Condi-Plus (2 ml/l)**

## Weak egg shells

Curative

**Vita-Min-DCP** for 21 days

- Group: 5 ml/l or 10 ml/kg feed
- Individually through the beak: 1 ml/kg body weight

Worming day 1, day 7 and day 21

- Group: **Vermi-Complete** (5 ml/l)
- Individually through the beak: **Endo-Ecto Plus Caps**: 1 capsule/kg body weight, or **Vermi-Complete** 1 ml/kg body weight

Prevention:

Check mixed faeces every 3-4 months:

- Make sure you also have mixed faeces from 2-3 days analysed for endoparasites - these may be responsible for deficiencies, and cause irregularities in eggshells

Vaccination RT+IBmulti+ND+EDS

- Viral infections and malformations near the fallopian tube



71

## Snot



**Mild infection + new animal**  
7-14 days  
**Respisor (3 ml/l)**

## Wry neck/stargazing



**Wry neck**  
14-21 days  
**Vit E + Se**

**Group:**  
2 ml/l  
**Individual (beak):**  
0.2-0.5 ml/kg body weight

## Exhibition or show



**General form**  
7 days before the show  
**Immuno-Plus (2 ml/l)**

**Yellower legs**  
14-21 days before the show  
**PURE Yellow 1-2 g/kg**

If problems persist, make an appointment with your vet

# Export prices

Please find the NeorniPharma export prices for private individuals and professionals below. These prices are **exclusive of VAT**. The VAT rates for the country of destination are applied when shipping. As always, professionals are VAT exempt.

PRODUCT	€ excl. VAT	PRODUCT	€ excl. VAT
Immuno Booster NEW (1 kg)	53.72	CU-For Liquid (250 ml)	12.40
Immuno-Plus (100 ml)	20.66	SilverFloX (1 l)	26.45
Immuno-Plus (250 ml)	45.45	Trichoryza Drops (50 ml)	14.05
Immuno-Plus (1 l)	147.93	Trichoryza Drops (250 ml)	48.76
Immuno Recup Caps (80 capsules)	19.83	Trichoryza Drops (1 l)	136.36
Super Start-Up (200 g)	20.66	Aminovital (100 ml)	15.70
Vit E + Se (100 ml)	14.05	Aminovital (250 ml)	28.93
Vit E + Se (250 ml)	28.93	Aminovital (1 l)	53.72
Hepa-Plus (100 ml)	16.53	Condi-Plus (100 ml)	12.40
Hepa-Plus (250 ml)	29.75	Condi-Plus (250 ml)	21.49
Hepa-Plus (1 l)	79.34	Condi-Plus (1 l)	68.60
Ready (100 ml)	18.18	Vita Energy (400 g)	24.79
Ready (250 ml)	39.67	Vita-Min-DCP (100 ml)	14.05
Ready (1 l)	119.01	Vita-Min-DCP (250 ml)	26.45
Dysbac Liquid (250 ml)	30.58	Vita-Min-DCP (1 l)	72.73
Feather Energy (400 ml)	15.70	Vita-Plus (100 ml)	14.88
Freeson Drops (30 ml)	14.05	Vita-Plus (250 ml)	23.14
PikStop (100 ml)	23.14	Vita-Plus (1 l)	53.72
Respicor (100 ml)	15.70		
Respicor (250 ml)	28.93	PURE	
Derma-Plus (50 ml)	28.93	Brewer's yeast (1 kg)	14.88
Exmite Spray (250 ml)	18.18	Dextrose (1 kg)	14.88
Exmite Spray (1 l)	57.02	Electrolyte (500 g)	14.88
Neorni Stabilizer (1 l)	16.53	Yellow (100 g)	12.40
Nositol (50 ml)	8.26	Yellow (500 g)	49.59
Probi-Plus (200 g)	14.88	Highly digestible protein (400 g)	20.66
SilverSpray (500 ml)	12.40	Red (100 g)	23.14
Skin Spray (200 ml)	14.05	Red (500 g)	70.25

# Index

Aminovital .....	42
Brewer's yeast .....	48
Condi-Plus .....	43
CU-For Liquid .....	38
Derma-Plus .....	28
Dextrose .....	48
Dosing card .....	16
Dysbacteriose .....	18
Dysbac Liquid .....	19
Electrolyte .....	48
ExmiteSpray .....	29
Feather Energy .....	22
Freeson Drops .....	23
Yellow .....	48
Hepa-Plus .....	14
Highly digestible protein .....	48
Immuno Booster NEW .....	7
Immuno-Plus .....	8
Immuno Recup Caps .....	9
Natural breeding promotion .....	6
Neorni Stabilizer .....	32
Nositol .....	33
PikStop .....	24
Probi-Plus .....	34
Ready .....	15
Red .....	48
Respicor .....	25
Schedules for pigeons .....	60-64
Schedules for chickens .....	70-71
Schedules for psittacidae .....	66-67
Schedules for songbirds .....	50-57
SilverFlox .....	39
SilverSpray .....	35
Skin Spray .....	36
Super Start-Up .....	10
Trichoryza Drops .....	40
Vita Energy .....	44
Vita-Min-DCP .....	45
Vita-Plus .....	46
Vit E + Se .....	11
Fertility therapy .....	12

EN

T +32 (0)11 96 91 02  
E [info@neornipharma.be](mailto:info@neornipharma.be)  
W [www.neornipharma.be](http://www.neornipharma.be)

**Desk & depot:**  
Beverlosesteenweg 129  
3583 Paal, Belgium

