

**SUPPLEMENTS**

**2026**



**NEORNI  
PHARMA**



# NEORNIPHARMA SUPPLEMENTS

## PREVENTIVE FORMULATIONS TAILORED TO YOUR BIRDS

This catalogue is a comprehensive guide of the NeorniPharma range aimed at **disease prevention through natural, highly concentrated supplements**. This range offers a safe and successful alternative to medicinal treatments and a long-term solution. By targeting annually recurring problems, NeorniPharma aims to build immunity and consequently stronger birds.

### Front desk address

Beverlosesteenweg 129  
3583 Paal, Belgium

### Phone number

+32 (0)11 96 91 02

### Additional dosages of powders

View alternative dosages on our  
'Powders and capsules' dosage card  
in this catalogue.



3 / 1

### Bird species in this catalogue



Pigeons



Songbirds



Fowl



Psittacidae



Birds of prey

All NeorniPharma products have unique formulas and are designed by veterinarians based on scientific research and practical experience.

## Breeding and immunity

| PRODUCT                         | DOSAGE                 | INDICATION                                 | PRICE (€) |
|---------------------------------|------------------------|--------------------------------------------|-----------|
| Immuno Booster NEW (1 kg)       | 20 g/kg egg food/ feed | All-in premix                              | 65        |
| Immuno-Plus (100 ml)            | 2 ml/l                 | Immunity, extra energy, breeding           | 25        |
| Immuno-Plus (250 ml)            | 2 ml/l                 | Immunity, extra energy, breeding           | 55        |
| Immuno-Plus (1 l)               | 2 ml/l                 | Immunity, extra energy, breeding           | 179       |
| Immuno Recup Caps (80 capsules) | 1 cap/0.5-1 kg BW      | Immunity, recovery, energy                 | 24        |
| Super Start-Up (200 g)          | 5 g/kg feed            | Start-up supplement for chicks, black spot | 25        |
| Vit E + Se (100 ml)             | 2 ml/l                 | Extra vitamin E and selenium               | 17        |
| Vit E + Se (250 ml)             | 2 ml/l                 | Extra vitamin E and selenium               | 35        |

## Liver, muscles, and condition

| PRODUCT            | DOSAGE | INDICATION                               | PRICE (€) |
|--------------------|--------|------------------------------------------|-----------|
| Hepa-Plus (100 ml) | 4 ml/l | Liver support                            | 20        |
| Hepa-Plus (250 ml) | 4 ml/l | Liver support                            | 36        |
| Hepa-Plus (1 l)    | 4 ml/l | Liver support                            | 79        |
| Ready (100 ml)     | 5 ml/l | Form, condition, and improved blood flow | 22        |
| Ready (250 ml)     | 5 ml/l | Form, condition, and improved blood flow | 48        |

## Megabacteria and dysbacteriosis

| PRODUCT                | DOSAGE | INDICATION                                     | PRICE (€) |
|------------------------|--------|------------------------------------------------|-----------|
| Dysbac Liquid (250 ml) | 2 ml/l | Crucible, proventriculus and intestinal health | 37        |

## Natural oils

| PRODUCT                 | DOSAGE       | INDICATION                     | PRICE (€) |
|-------------------------|--------------|--------------------------------|-----------|
| Feather Energy (400 ml) | 5 ml/kg feed | Fly oil                        | 19        |
| PikStop (100 ml)        | 1 ml/l       | Pecking                        | 28        |
| Respicor (100 ml)       | 3 ml/l       | Respiratory tract, heat stress | 19        |
| Respicor (250 ml)       | 3 ml/l       | Respiratory tract, heat stress | 35        |

## Probiotics and disinfection

| PRODUCT                 | DOSAGE         | INDICATION                   | PRICE (€) |
|-------------------------|----------------|------------------------------|-----------|
| Neorni Stabilizer (1 l) | 5-10 ml/100 ml | Probiotics - environment     | 20        |
| Probi-Plus (150 g)      | 3 g/l          | Probiotics and prebiotics    | 18        |
| SilverSpray (500 ml)    | Spray:         | Disinfectant spray           | 15        |
| Skin Spray (200 ml)     | Spray:         | Probiotics - skin infections | 17        |



## Acidification and drinking water disinfection

| PRODUCT                   | DOSAGE             | INDICATION                      | PRICE (€) |
|---------------------------|--------------------|---------------------------------|-----------|
| CU-For Liquid (250 ml)    | 1-2 ml/l           | Acidification of drinking water | 15        |
| SilverFloX (1 l)          | 2 ml/l             | Drinking water disinfection     | 32        |
| Trichoryza Drops (50 ml)  | 1 drop per nostril | Throat elixir                   | 17        |
| Trichoryza Drops (250 ml) | 10 ml/kg           | Throat elixir                   | 59        |
| Trichoryza Drops (1 l)    | 10 ml/kg           | Throat elixir                   | 165       |

## Vitamins, amino acids, electrolytes, minerals, and trace elements

| PRODUCT               | DOSAGE       | INDICATION                                                | PRICE (€) |
|-----------------------|--------------|-----------------------------------------------------------|-----------|
| Aminovital (100 ml)   | 2 ml/l       | Amino acids, electrolytes, vitamins                       | 19        |
| Aminovital (250 ml)   | 2 ml/l       | Amino acids, electrolytes, vitamins                       | 35        |
| Aminovital (1 l)      | 2 ml/l       | Amino acids, electrolytes, vitamins                       | 65        |
| Condi-Plus (250 ml)   | 2 ml/l       | Moulting supplement                                       | 26        |
| Vita Energy (400 g)   | 10 g/kg feed | Conditioning powder – vitamins, minerals, and amino acids | 30        |
| Vita-Min-DCP (100 ml) | 2 ml/l       | Vit D, calcium, phosphorus, and other minerals            | 20        |
| Vita-Min-DCP (250 ml) | 2 ml/l       | Vit D, calcium, phosphorus, and other minerals            | 35        |
| Vita-Min-DCP (1 l)    | 2 ml/l       | Vit D, calcium, phosphorus, and other minerals            | 70        |
| Vita-Plus (100 ml)    | 2 ml/l       | Water-soluble multivitamins                               | 18        |
| Vita-Plus (250 ml)    | 2 ml/l       | Water-soluble multivitamins                               | 28        |
| Vita-Plus (1 l)       | 2 ml/l       | Water-soluble multivitamins                               | 65        |

## PURE

| PRODUCT                           | DOSAGE          | INDICATION                                   | PRICE (€) |
|-----------------------------------|-----------------|----------------------------------------------|-----------|
| Brewer's yeast (1 kg)             | 15-30 g/kg feed | Amino acids, minerals, trace elements, Vit B | 18        |
| Dextrose (1 kg)                   | 15-30 g/l       | Energy, flavour                              | 18        |
| Electrolyte (500 g)               | 15-30 g/l       | Hydration, stabilisation, balance            | 18        |
| Yellow (100 g)                    | 1-2 g/kg feed   | Colour                                       | 15        |
| Yellow (500 g)                    | 1-2 g/kg feed   | Colour                                       | 60        |
| Highly digestible protein (400 g) | 5-15 g/kg feed  | Muscle growth, extra amino acids/proteins    | 25        |
| Red (100 g)                       | 1 g/kg feed     | Colour                                       | 28        |
| Red (500 g)                       | 1 g/kg feed     | Colour                                       | 85        |

# NEORNI

NeorniPharma is part of Neorni, a team of veterinarians dedicated specifically to avian medicine. Neorni offers various services to ensure the best care for birds.

Follow us  
on Facebook



[www.neorni.be](http://www.neorni.be)  
[www.neornilab.be](http://www.neornilab.be)



## NeorniVet Huldenberg

[huldenberg@neornivet.be](mailto:huldenberg@neornivet.be)  
OPENING 2026



NEORNI<sup>VET</sup>

## Veterinary practice

NeorniVet Paal  
NeorniVet Huldenberg



NEORNI<sup>LAB</sup>

## Laboratory

Gender • Performance genes  
Kinship • Faecal analysis  
Disease testing • Bacteriology  
Species determination  
Autopsy • Colour mutations

NEORNI



PHARMA


BY VETS FOR BIRDS

## Product range



## Neorni (all services)

Beverlosesteenvweg 129  
3583 Paal, Belgium  
T +32 (0)11 96 91 02



# BREEDING AND IMMUNITY



# Natural breeding boost

## Basic breeding

The basic breeding schedule is the essence of successful breeding. Several field trials have shown that the birds were better prepared for breeding, which was reflected in more nests and more young per nest.



### Immuno-Plus

2 ml/l or 4 ml/kg

**Prior to breeding:**  
daily for 2-6 weeks until  
the first clutch

Breeding urge +++

Polypeptides

Stimulation of egg  
cell and sperm cell  
production



### Immuno Booster NEW

20 g/kg egg food/feed

**During breeding:**  
daily from the first clutch un-  
til the young leave the nest

Breeding urge +

Polypeptides and Premix  
breeding and rearing

Maintenance of egg cell and  
sperm cell production



### Need advice?

Your Neorni  
veterinarian will be  
happy to help



### Vit E + Se

2 ml/l or 4 ml/kg

Daily for 2-4 weeks before  
breeding until the first  
clutch

Fertility +++

Linseed oil, arginine,  
spirulina, carnitine, zinc, etc

Extra high dose of  
Vit E + Se

6

## Extra breeding (fertilisation)

Fertilisation problems occurring upon application of the basic schedules often have a nutritional or infectious cause.

►► nutritionally speaking, a **Vit E + Se** supplement would certainly be beneficial.

►► In addition, certain germs with an important influence on fertilisation have been isolated in recent years. Faeces culture before breeding can be carried out easily and quickly at NeorniVet veterinary practice(± 24h).

►► In specific situations/species, extra added value can be achieved with the use of specific supplements such as **Vita-Plus, Ready, Hepa-Plus ...**

# IMMUNO BOOSTER NEW



## All-in premix for young birds



### Indication

- Supports breeding results and maintains the condition of your birds
- Supply of extra energy
- Supports the immune system
- Uses probiotica to help build up intestinal flora
- Good development of the skeletal system
- Ideal for animals that are required to deliver top performances

### Administration 20 g/kg

- 1 large scoop (20 g)/kg egg food/feed or 2 small scoops (2 g)/100 g egg food/feed
- During breeding season: add daily to egg food/feed
- Extra energy and accommodating deficits: 1-2 days per week throughout the year

*Corn gluten, potato protein, potato protein fermentate, yeast and parts of it, chicory root fibre, spirulina, yeast, trimagnesium citrate, monoglycerides, diglycerides and triglycerides of fatty acids (c4), tricalcium citrate, L-carnitine, betaine, vitamin A, vitamin D3, vitamin E, vitamin C vitamin B1, vitamin B2, calcium D-pantothenate, vitamin B6, vitamin B12, vitamin K, niacinamide, folic acid, choline, clinoptilolite, copper, iron, zinc, manganese, potassium iodide, selenium, cobalt, lysine, methionine, glycine, echinacea purpurea, curcuma, lactobacillus acidophilus*





## Immunity, extra energy, breeding



100 ml €25

250 ml €55

1 l €179

"Vogel-Viagra"

8

### Indication

- Supports breeding results and maintains the condition of your birds
- Supply of extra energy
- Good appetite and feather patterns
- To be used before/after intense exertion/competition
- Supports the development and growth of young animals

### Administration 2 ml/l drinking water

- Breeding: daily from 2-6 weeks before breeding. Stop administration from first clutch.
- Show/competition: starting 5 days before show/competition, daily 2 ml/l drinking water or 2 ml/kg body weight directly into the beak
- Immunosuppression and disease: 2 ml/l drinking water for 14 days or 2 ml/kg body weight directly into the beak until recovery
- Viral infections (circovirus, bornavirus, etc.): for 4-8 weeks to support and promote immunity

# IMMUNO RECUP CAPS



## Immunity, recovery, energy



80 capsules

€24

### Indication

- Stimulates recovery after illness/exertion
- Supply of extra energy
- Supports the immune system
- Ideal for top performance

### Administration

- Recovery/immunity boost: 1 capsule/0.5-1 kg orally for 5-10 days
- Extra energy: 1 capsule/0.5-1 kg orally
- Pigeon: day of return home, 1 capsule/pigeon
- To support breeding: for 14 days before mating until laying 1 capsule/0.5-1 kg
- *Immuno Recup Caps* can be administered directly or with the food. When administered via food, the full dose should be consumed.

*Maize gluten, potato protein, potato protein fermentative, yeast and parts of it, chicory root fibre, spirulina, yeast, trimagnesium citrate, monoglycerides, diglycerides and triglycerides of fatty acids (c4), tricalcium citrate, L-carnitine, Vit A, Vit D3, Vit E, Vit C, Vit B1, Vit B2, calcium D-pantothenate, Vit B6, Vit B12, Vit K, niacinamide, folic acid, choline, clinoptilolite, copper, iron, zinc, manganese, potassium iodide, selenium, cobalt, lysine, methionine, glycine, echinaceae purpureae, curcuma, lactobacillus acidophilus*

200 g €25



- Administer daily from the first clutch until the young leave the nest



# VIT E + SE



## Extra vitamin E + selenium



100 ml €17

250 ml €35

+ spirulina  
+ zinc  
+ linseed oil

### Indication

- For animals with a deficiency of vitamin E selenium
- Recommended for nervous symptoms

### Administration

- 2 ml/l drinking water or 1 ml/kg body weight directly into the beak
- Recommended in case of nervous symptoms: for 7-10 days




#### **Fertility +++**

Daily 2 ml/l or 4 ml/kg feed daily for 2-4 weeks, prior to breeding until the first clutch

# Fertility therapy

Individual therapy (cocks ♂ and hens ♀)





# LIVER, MUSCLES, AND CONDITION



## Liver support



14

### Indication

- Supports liver function
- Relieves stress on the liver

### Administration

- Curative: 4 ml/l drinking water for 5-14 days
- Prevention: 2 ml/l drinking water, 2 to 3 days per week



Can be administered daily to support the liver during administration of medication or dye



## Form, condition, and improved blood flow



100 ml €22

250 ml €48

*Additional  
iron, iodine,  
L-carnitine,  
and Vit B12*

### Indication

- Condition booster: improves flight performance
- Stimulates fat metabolism and endurance
- Supports blood circulation


### Administration 5 ml/l drinking water

- **Condition boost:**
  - For exhibition/flying season: for 2 weeks
  - For important flight: for 2-5 days
- **Fat birds:** for 3-5 days a week until the result is achieved
- **Anaemia:** 7-10 days
- Adjusted dosage songbirds <250 g body weight (canary, siskin, etc.): 1-2 ml/l

*Glycerol, L-carnitine, beetroot juice, potassium iodide, iron glycinate, vitamin B12, magnesium citrate sodium chloride, magnesium chloride, calcium chloride, glucose, vitamin E, selenium, zinc, copper, manganese, vitamin A, vitamin B1, vitamin B2, vitamin B5, vitamin B6, vitamin PP (niacinamide), vitamin D3, vitamin C, blend of natural aromatic components*

## Notes

This image shows a full page of primary-ruled paper. It features multiple sets of horizontal lines designed to guide handwriting. Each set consists of three lines: a solid top line, a dashed middle line, and a solid bottom line. These sets are repeated vertically down the entire page, providing ample space for practicing letter formation and alignment. The paper is otherwise blank, with no margins or additional markings.



# MEGABACTERIA AND DYSBACTERIOSIS

# Dysbacteriosis

crop  
proventriculus  
intestines



**Dysbac Liquid**  
prevention of  
dysbacteriosis

**7-in-1**  
Intestinal health  
on 7 fronts

**Dysbac Liquid** is an improved and extended version of **CU-For Liquid** with more focus on **intestinal health** and **general immunity**

18



2 ml/l

+

+++

+++

+++

+++

+++

**CROP, PROVENTRICULUS AND  
INTESTINAL HEALTH, PREVENTION  
OF DYSBACTERIOSIS**

►► **Short-chain fatty acids:**  
formic acid, propionic acid, citric acid

►► **Medium-chain fatty acids:**  
butyric acid, monoglycerides, diglycerides and triglycerides of fatty acids (C8-C10)

►► **Aromatic substances:**  
oregano, echinaceae purpureae, ginger, cinnamon, black pepper, garlic, and boswellia

►► **Vitamin C**

►► **Chelated copper and zinc**

►► **Extra fibre**

►► **Fish oil**

**DOSAGE**

ACIDIFICATION OF DRINKING WATER (pH)

CROP HEALTH

POVENTRICULUS HEALTH

INTESTINAL HEALTH

GENERAL IMMUNITY

**PURPOSE**

1-2 ml/l

+++

+++

++

-

-

**WATER ACIDIFICATION,  
WATER QUALITY**

►► **Short-chain fatty acids:**  
ammonium formate, acetic acid, lactic acid, formic acid, propionic acid

►► **Copper and zinc sulphates**



**Need advice?**

Your Neorni  
veterinarian will be  
happy to help





## Crop, proventriculus and intestinal health



### Indication

- Forms a protective film over the intestinal mucosa
- Increases the animal's immunity
- Helps to maintain the natural balance of intestinal flora
- To be used in cases of watery faeces

### Administration 2 ml/l drinking water

- Prevention: 1 to 2 days per week
- In case of digestion problems: for 7-14 days



### Prevention of megabacteriosis and dysbacteriosis

- 1-2 days a week 2 ml/l drinking water
- May be combined with ACS

# Trichoryza Drops

The best way to tackle  
trichomoniasis  
**without losing form**

**FLYING SEASON**  
10 ml/  
kg feed  
1 day/week, day after returning home

**NEW**  
1 litre



**Proven effectiveness  
against resistant  
trichomonas!**

---



# NATURAL OILS



## Flight oil, source of energy from lipids



22

### Indication

- Source of energy from lipids
- Feather quality, optimal feathering
- Extra vitamins A, D, and E
- Hoge source of omega 3-6-9 fatty acids

### Administration 5 ml/kg feed

- Moulting: 2 days per week
- Breeding: 2-3 days per week
- Flying season:
  - Departure: for 2-5 days before basketing, in combination with *Vita Energy*
  - Homecoming: for 2 days, in combination with *Immuno Booster NEW*



Can be used to improve the absorption of *Immuno Booster NEW* and *Vita Energy* with the feed

Cold-pressed oils: linseed oil, cod liver oil, rapeseed oil, salmon oil, lecithin, maize germ oil, wheat germ oil, sunflower oil, hemp seed oil (including vitamins A, D and E)

## Notes

This image shows a full page of primary-ruled paper. It features multiple sets of horizontal lines designed to guide handwriting. Each set consists of three lines: a solid top line, a dashed middle line, and a solid bottom line. These sets are repeated vertically down the entire page, providing ample space for practicing letter formation and alignment. The paper is otherwise blank, with no margins or additional markings.



## Pecking, restlessness



NEW

100 ml €28



24

### Indication

- Pecking in birds and poultry

### Administration

- 1 ml/l drinking water for 4-6 weeks



#### Feather picking/pecking

Administer simultaneously with *Immuno-Plus* for the best results



## Respiratory tract, heat stress



100 ml €19

250 ml €35

formerly  
**Herbi+E**

### Indication

- Natural solution to support respiratory problems
- Suppression of pain/inflammation in case of respiratory problems
- Stuck slime
- Heat stress

**Administration** 3 ml/l drinking water or 0.5 ml/kg body weight directly into the beak

- Preventive: when purchasing new birds: for 10-14 days
- Curative: in mild to moderate respiratory problems, for 10-14 days
- Heat stress: 3-5 ml/l



**Additional support for racing pigeons in hot weather** 3 ml/l of drinking water, in combination with *Aminovital* on the basketing day

2026

# neofalco


products for falcons



more info: [info@neornipharma.be](mailto:info@neornipharma.be)



---



# PROBIOTICS AND DISINFECTION

# NEORNI STABILIZER



## Probiotic ambient spray



1 | €20

28

### Indication

- Reduction of pathogenic load in your birds' surroundings
- Forms a stable and healthy environment by occupying it with safe, probiotic bacteria

### Administration

- To be used diluted: 5-10% (5-10 ml/100 ml) in lukewarm water
- Mist the entire area to be stabilised
- First week, spray the loft daily (may be done in the presence of birds, including nestlings).
- Subsequently, spray twice a week to maintain probiotic protection
- The product diluted with water has a maximum shelf life of 5 days.



Can be nebulised in the presence of the birds/pigeons, especially during important or stressful periods when the bacterial presence in the environment rises

## Notes

[illegible]

150 g €18



- Increases the beneficial flora in the crop and the intestines
- Ensures improved digestion and supports the initiation of the crop and intestinal flora in chicks
- After a course of antibiotics

- Preventive: for 2 days/week
- Wet droppings or after antibiotic treatment: for 5-7 days
- Refresh daily
- May be combined with other supplements (vitamins)



Change drinking water twice daily or administer through the feed



## Disinfectant spray



### Indication

- Broad-spectrum disinfectant
- Bactericide - Fungicide - Virucide

### Administration

- Ready-to-use disinfectant spray
- Suitable for lofts, cages, equipment, and eggs
- Indicated when cleaning soiled plumage (faeces, blood)



Safe to apply in the presence of your birds



## Probiotic spray for skin infections



200 ml €17

32

### Indication

- Suitable for birds with sensitive skin
- To be used externally for care of the skin
- Reduces the formation of scaly skin

### Administration


- Mist the area to be stabilised



#### **Powder comb/flavus/white flakes**

Spray the comb or infected area daily. Not harmful to eyes or respiratory tract.

---



# ACIDIFICATION AND DRINKING WATER DISINFECTION



## Acidification of drinking water



250 ml €15

34

### Indication

- Acidification of drinking water, lowering pH

### Administration 1-2 ml/l drinking water

- Prevention: 1 to 2 days per week



Exceptionally concentrated formula, very economical to use





## Drinking water disinfection



NEW

1 | €32

### Indication

- Water disinfection
- Prevention of harmful bacteria, viruses, yeasts and moulds
- Immunity support

### Administration

- 2 ml/l drinking water



Longer and better action due to silver addition



## Throat elixir



36

### Indication

- Preventive approach to trichomoniasis
- Supports throat and crop health
- Positive effect on the upper respiratory tract
- Plays a role in resistance to disease

### Administration

- The day after returning home:
  - Recommended administration: 10 ml/kg feed
- **Always make fresh before each feeding. Most effective up to 2 hours after mixing.**
- 5 ml/l drinking water
- 1 drop per nostril
- Prevention: 1 day per week.
- At high pressure: 2 days per week
- Caution: stains clothing and hands



Proven efficacy in resistant trichomoniasis cases:  
20 ml/kg feed for 5 days

---



# VITAMINS, AMINO ACIDS, ELECTROLYTES, MINERALS, AND TRACE ELEMENTS





## Amino acids, electrolytes, and vitamins



100 ml €19

250 ml €35 1

| €65

Extra vitamin E,  
selenium and BCAA  
amino acids

38

### Indication

- All essential amino acids crucial for proper moulting/feather development/pigmentation
- Stimulates recovery/repair
- Muscle building and recovery

### Administration 2 ml/l drinking water

- Racing pigeons: 5 ml/l during the racing season
- Throughout the entire year: 1 to 2 days per week
- Throughout the entire moulting period: 3 days/ week. May be combined with *Condi-Plus*.



**Additional support for racing pigeons in warm weather** 5 ml/l drinking water, in combination with *Respicor* on the basketing day

*Dextrose, glycerol, sodium chloride, prebiotic fibre, trimagnesium citrate, beetroot juice, potassium chloride, sucrose, calcium chloride, tricalcium citrate, lysine, methionine, threonine, tryptophan, histidine, carnitine, betaine, choline, vitamin B1, vitamin B2, niacinamide, calcium D-pantothenate, vitamin B6, vitamin B12, folic acid, biotin, vitamin C, vitamin E, selenium, copper, zinc, manganese, mix of natural aromatic components, L-isoleucine, L-valine, L-leucine*



## Moulting supplement



250 ml €26

### Indication

- Supports bird health during moulting
- Supports the metabolism
- Supply of extra energy
- Increases appetite

### Administration 2 ml/l drinking water

- Throughout the entire moulting period: 3 days/ week. May be combined with *Aminovital*.



## Conditioning powder – Vitamins, minerals, and amino acids



40

### Indication

- Condition booster and extra energy
- Extra vitamins, minerals and amino acids
- Increases endurance
- Breeding stimulation

### Administration

- Recommended administration: 10 g/kg feed
- Administration via drinking water: 4 g/l



- General: Throughout the year, 1-2 days/week.  
In case of poor/impaired condition: for 14 days.
- Breeding stimulation: 2-4 weeks prior to breeding, twice a week to daily
- Pigeon: Medium distance/long distance, 2-5 days before basketing

*Dextrose, maltodextrin, soya protein concentrate, calcium carbonate, yeasts and parts thereof, trimagnesium citrate, monosodium phosphate, sodium bicarbonate, sodium chloride, potassium chloride, trisodium phosphate, tricalcium citrate, vitamin A, vitamin D3, vitamin E, vitamin K, vitamin B1, vitamin B2, vitamin B3, vitamin B5, vitamin B6, vitamin B11, vitamin B12, biotin, vitamin C, choline chloride, iron, copper, zinc, manganese, potassium iodide, selenium, lysine, methionine, tryptophan, threonine, mixture of natural aromatic components*



## Liquid source of vitamin D, calcium, phosphorus, and other minerals



100 ml €20

250 ml €35

1 l €70

Extra calcium  
and  
phosphorus

### Indication

- For birds with a deficiency in Vitamin D3, calcium, phosphorus and other minerals
- Supports bone formation and development in young animals
- Supports egg shell formation
- Poor egg shell quality/shell-less eggs

### Administration 2 ml/l drinking water

- Prevention: 2 ml/l
  - 2 weeks prior to breeding until the eggs are laid: 2-3 days per week
  - For the first 6 weeks of life: 2-3 days per week
- Curative: 5 ml/l for 14 days
- Support for fractures: for 6 weeks, 5 ml/l drinking water or 0.5 ml/kg body weight directly into the beak



Change drinking water twice daily to avoid sedimentation, or administer via the feed (preventive: 4 ml/kg egg food/feed - curative: 10 ml/kg egg food/feed)



## Water-soluble multivitamins



42

### Indication

- Supplement containing all essential vitamins
- Overall daily health
- Prevention of chronic vitamin A deficiency

### Administration 2 ml/l drinking water

- Weekly: for 1 to 2 days per week
- After illness/recuperation: for 3 to 5 consecutive days



For parrots and parakeets, we strongly recommend administering 1-2 days per week *Vita-Plus* via the drinking water to prevent vitamin A deficiency



High concentration of vitamin K helps prevent or stop bleeding



---



# PURE PURE

# BREWER'S YEAST



## Indication

- Amino acids, minerals and trace elements
- Vitamin B

**Administration** 15-30 g/kg feed

1 kg €18

# DEXTROSE



## Indication

- Additional energy source
- Flavouring medicated water

**Administration** 15-30 g/l drinking water

1 kg €18

# ELECTROLYTE



## Indication

- Hydration in case of dehydration
- Stabilisation of water and electrolyte balance

**Administration** 15-30 g/l drinking water

500 g €18

44

# YELLOW



## Indication


- Colour

**Administration** 1-2 g/kg feed

100 g €15

500 g €60

# HIGHLY DIGESTIBLE PROTEIN



## Indication

- Muscle growth
- Additional amino acids/proteins

**Administration** 5-15 g/kg feed

400 g €25

# RED



## Indication

- Colour

**Administration** 1 g/kg feed

100 g €28


500 g €85



# SCHEDULES FOR SONGBIRDS

# Annual Euro and Canary schedule

## Timeline



### Preparation for breeding

- Breeding boost
- Liver support
- Black spot prevention

### First clutch until weaning

- Optimal start and mushy chick disease prevention
- Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi

### Weaned young

- Prevention coccidiosis/atoxoplasmosis, megabacteriosis and dysbacteriosis
- Limiting pressure with immunity build-up





# Annual Euro & Canary schedule

*The annual schedule is a general guide for preparation, breeding, hatching, weaning and moulting. Talk to your veterinarian about the specific requirements for your birds and devise a tailored schedule together.*

## Preparation from 6-8 weeks prior to breeding

1. A healthy environment is the best start! (see Virkon™S or SilverSpray)
2. Prevention of atoxoplasmosis/coccidiosis + megabacteriosis/dysbacteriosis
  - **Veterinarian:** bird check-up + faecal analysis 6-8 weeks prior to breeding
  - Only use courses of treatment in case of problems:
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 3 days Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
3. For 6 weeks prior to breeding
  - **Immuno-Plus (2 ml/l) daily**
    - Breeding boost; birds will be in top condition
  - 1-2 days per week **Hepa-Plus (4 ml/l)**
    - Liver support
  - As required: Vita-Plus, Vita Energy, Ready, Vit E + Se, Vita-Min-DCP
    - These are suitable for use with Immuno-Plus
  - Extra: black spot prevention
    - For 7 days **Gould Mix (5 g/l)** during breeding preparation
4. Prevention of ectoparasites
  - Haematophagous mites
    - **Exzolt:** 1 ml/l for 1 day, repeat on days 7 and 14
      - Does not affect fertility and nestlings/young
      - Can be repeated every 3 months
    - Extra prevention of haematophagous mites
      - Before occupation, treat loft with:
        - **Derma-Plus**
  - Feather mites
    - **Vermi-Complete (1 ml/l bath water)**

47

## Breeding: First clutch until weaning

1. **Immuno Booster NEW (2 g/100 g or 20 g/kg egg food/feed)**
  - All-in premix for young birds
2. First 6 weeks
  - Option 1: **Super Start-Up (5 g/kg egg food/feed)**
    -  Natural suppression and prevention of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
  - Option 2: **Super Start-Up Blue (20 g/kg egg food/feed)**
    -  Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi



3. Start from 2 days before hatching until 5 days after hatching **Coli-Mix (5 g/l)**
  - Prevention of mushy chick disease (Colibacillosis)
  - Alternative to Coli-Mix: **Gould Mix (5 g/l)**

### Breeding: weaned young

1. A healthy environment is the best start!

- **Virkon™S or SilverSpray**
  - Environment free of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and moulds
- **Neorni Stabilizer:** Probiotic ambient spray
  - Prevention by reducing bacterial presence in the environment

2. Preventive cure

- Third week after weaning
  - **Increased susceptibility to atoxoplasmosis/megabacteriosis**
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 3 days Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)

3. Then throughout the year (from weaning to breeding period)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis and dysbacteriosis

48

### Moulting period

Throughout moulting **Condi-Plus + Aminovital:**

- Condi-Plus (2 ml/l): 3 days/week
- Aminovital (2 ml/l): 3 days/week
- May be administered simultaneously

**Intermediate season** Throughout the year (from weaning to breeding)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis and dysbacteriosis
- As required: Vita-Plus, Vita Energy, Ready, Vit E + Se, Vita-Min-DCP

At NeorniPharma, we always strive for the best long-term solution. We are constantly looking for **alternatives to anti-microbials** to limit the risk of resistance. The goal is to give our birds **as few preventive treatments with antimicrobial agents as possible**. Clinical research and any additional tests are crucial for this. After the diagnosis, an appropriate treatment can be administered. **Antibiotics are not harmless**. Consult your vet for advice about their correct and responsible use. **Autopsy:** don't simply throw away a bird that has died! The cadaver can provide a lot of information to ensure appropriate treatment can be sought. The cadavers are best kept cool (fridge) and analysed within 24 hours (max. 72 hours).



# Preparation for breeding



## Need advice?

Your Neorni veterinarian will be happy to help

| Immuno-Plus                                       | Hepa-Plus                                      |
|---------------------------------------------------|------------------------------------------------|
| 2 ml/l                                            | 4 ml/l                                         |
| 6 weeks prior to mating until the first clutch    | 6 weeks prior to mating until the first clutch |
| Daily                                             | 1-2 days/week                                  |
| Stimulation of sperm cell and egg cell production | Supports the liver                             |

49



| Vit E + Se                                                                              | Vita-Min-DCP                                          | Ready                                          | Vita-Plus/Vita Energy                                                      |
|-----------------------------------------------------------------------------------------|-------------------------------------------------------|------------------------------------------------|----------------------------------------------------------------------------|
| 2 ml/l                                                                                  | 2 ml/l                                                | 3-5 ml/l                                       | 2 ml/l or 10 g/kg                                                          |
| 2-4 weeks prior to mating until the first clutch                                        | 2 weeks prior to mating until the first clutch        | 6 weeks prior to mating until the first clutch | A: Throughout the year:<br>B: 2 weeks before mating until the first clutch |
| Daily                                                                                   | 2 days/week                                           | 1-2 days/week                                  | A: 1-2 days/week<br>B: Daily                                               |
| Extra-high doses of Vit E + Se, linseed oil, arginine, spirulina, carnitine, zinc, etc. | Extra calcium, phosphorus, Vit D3, and trace minerals | Converts fat into usable energy for the bird   | Extra vitamins                                                             |

**FERTILISATION**

**EGGSHELL**

**CONDITION**

**VITAMINS**

▶▶▶▶ All products can be used in combination





# Breeding: hatching to weaning



50

**Coli-Mix** 

5 g/l or 10 g/kg egg food/feed

2 days before to 5 days after hatching

No impact on fertility

Broad-spectrum mix

**Prevention of mushy chick disease and Bengalese finch disease**

**Immuno Booster NEW**

20 g/kg egg food/feed

Daily from the first clutch until the young leave the nest

Positive effect on fertility

All-in premix

**Maintain oocyte and sperm production**

**Super Start-Up** 

5 g/kg egg food/feed

**Natural suppression and prevention of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi**

**OR**

**Super Start-Up Blue** 

20 g/kg egg food/feed

**Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi**

**Immuno Booster NEW** contains all essential components for **rearing young birds**. It is a unique **comprehensive** premix to mix under egg food/feed, providing all necessary nutrients in **exceptionally high concentrations**.

**Immuno Booster NEW Polypeptides** Keep the birds in form during breeding. **Saccharomyces cerevisiae** Source of protein, amino acids, and nucleotides. **Chicory root fibre** Prebiotic fibre: balanced digestion. **Spirulina** Source of protein, fats, carbohydrates, vitamins, and minerals. **Electrolytes** Fluid balance. **Fatty acids** Energy, balanced digestion. **All water-soluble and fat-soluble vitamins** General immunity, antioxidants, growth, energy release. **All amino acids** Building muscle and feather cover. **L-carnitine** Supports fat digestion and muscle building. **Choline** Liver support. **Minerals** (cobalt, iron, copper, zinc, manganese, iodine, selenium) General immunity, antioxidants, growth, energy release. **Clinoptoplite** General health and intestinal health. **Probiotics** Intestinal flora stabiliser. **Turmeric and echinacea** General immunity and health.





# Breeding: weaned young

| COCCIDIOSIS AND ATOXOPLASMOSIS               |  | MEGABACTERIA AND DYSBACTERIOSIS                  |  |
|----------------------------------------------|--|--------------------------------------------------|--|
| ACS                                          |  | Dysbac Liquid                                    |  |
| 0.5 ml/l                                     |  | 2 ml/l                                           |  |
| From weaning until next breeding time        |  | From weaning until next breeding time            |  |
| 3-5 days/month<br>If required: twice monthly |  | 1-2 days/week                                    |  |
| COCCIDIOSIS AND ATOXOPLASMOSIS               |  | MEGABACTERIOSIS, DYSBACTERIOSIS, AND FLAGELLATES |  |



## Extra support throughout the year

| SUPPORTS THE LIVER                             |  |
|------------------------------------------------|--|
| Hepa-Plus                                      |  |
| 2 ml/l                                         |  |
| Throughout the year                            |  |
| 1-2 days/week<br>When adding dye: 2 ml/l daily |  |
| LIVER SUPPORT                                  |  |



| GENERAL SUPPORT         |  |
|-------------------------|--|
| Vita-Plus / Vita Energy |  |
| 2 ml/l or 10 g/kg       |  |
| Throughout the year     |  |
| 1-2 days/week           |  |
| EXTRA VITAMINS          |  |



51



# Annual Exotics schedule

*The annual schedule is a general guide for preparation, breeding, hatching, weaning and moulting. Talk to your veterinarian about the specific requirements for your birds and devise a tailored schedule together.*

## Preparation from 6-8 weeks prior to breeding

1. A healthy environment is the best start! (see Virkon™S or SilverSpray)
2. Prevention of flagellates (cochlosomosis) + campylobacteriosis
  - **Veterinarian:** bird check-up + faecal analysis 6-8 weeks prior to breeding
  - Only use courses of treatment in case of problems:
    - For 7 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)
3. For 6 weeks prior to breeding
  - **Immuno-Plus (2 ml/l) daily**
    - Breeding boost; birds will be in top condition
  - 1-2 days per week **Hepa-Plus (4 ml/l)**
    - Liver support
  - As required: Vita-Plus, Vita Energy, Ready, Vit E + Se, Vita-Min-DCP
    - These are suitable for use with Immuno-Plus
  - Prevention of cochlosomosis + campylobacteriosis
    - For 7 days **Gould Mix (5 g/l)** during breeding preparation
4. Prevention of ectoparasites
  - Haematophagous mites
    - **Exzolt:** 1 ml/l for 1 day, repeat on days 7 and 14
      - Does not affect fertility and nestlings/young
      - Can be repeated every 3 months
    - Extra prevention of haematophagous mites
      - The day prior to occupation, treat loft with:
        - **Derma-Plus**
  - Tracheal mites
    - **Vermi-Complete:** 2.5-5 ml/l for 1 day, repeat after 3 weeks
  - Feather mites
    - **Vermi-Complete (1 ml/l bath water)**

## Breeding: First clutch until weaning

1. **Immuno Booster NEW (2 g/100 g or 20 g/kg egg food/feed)**
  - All-in premix for young birds
2. First 6 weeks
  - Option 1: **Super Start-Up (5 g/kg egg food/feed)**
    -  Natural suppression and prevention of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
  - Option 2: **Super Start-Up Blue (20 g/kg egg food/feed)**
    -  Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
3. Start from 2 days before hatching until 5 days after hatching **Coli-Mix (5 g/l)**
  - Prevention of mushy chick disease (Colibacillosis)
  - Alternative to Coli-Mix: **Gould Mix (5 g/l)**



#### 4. For problems in the first 6-8 weeks of life

- **Gould Mix (5 g/l)**
  - For the treatment of cochlosomosis (Bengalese finch disease), yeasts and bacterial diarrhoea

### Breeding: weaned young

#### 1. A healthy environment is the best start!

- **Virkon™S or SilverSpray**
  - Environment free of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and moulds
- **Neorni Stabilizer:** Probiotic ambient spray
  - Prevention by reducing bacterial presence in the environment

#### 2. Preventive cure

- Third week after weaning
  - **Increased susceptibility to coccidiosis/cochlosomosis/campylobacter**
    - 5 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)
    - 3 days Hepa-Plus (4 ml/l)
    - 5 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)

#### 3. Then throughout the year (from weaning to breeding period)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis and dysbacteriosis
  - Prevention of flagellates (cochlosomosis)

53

### Moulting period

Throughout moulting **Condi-Plus + Aminovital:**

- Condi-Plus (2 ml/l): 3 days/week
- Aminovital (2 ml/l): 3 days/week
- May be administered simultaneously

### Mid-season

Throughout the year (from weaning to breeding)

- **ACS 3-5 days (0.5 ml/l) 1-2 times per month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis and dysbacteriosis
  - Prevention of flagellates (cochlosomosis)
- As required: Vita-Plus, Vita Energy, Ready, Vit E + Se, Vita-Min-DCP

At NeorniPharma, we always strive for the best long-term solution. We are constantly looking for **alternatives to anti-microbials** to limit the risk of resistance. The goal is to give our birds **as few preventive treatments with antimicrobial agents as possible**. Clinical research and any additional tests are crucial for this. After the diagnosis, an appropriate treatment can be administered. **Antibiotics are not harmless**. Consult your vet for advice about their correct and responsible use. **Autopsy:** don't simply throw away a bird that has died! The cadaver can provide a lot of information to ensure appropriate treatment can be sought. The cadavers are best kept cool (fridge) and analysed within 24 hours (max. 72 hours).



# Vaccination schedule for pigeons

3-4 WEEKS INTERVAL

## Male

|                           |                                             |
|---------------------------|---------------------------------------------|
| Age 6 weeks               | <b>Rota-paramyxo</b>                        |
| Age 8 weeks               | Salmonella/paratyphoid herpes               |
| Age 10 weeks              | <b>Smallpox</b> (injection/follicle method) |
| 3-4 weeks before training | <b>Rota-paramyxo BOOSTER</b>                |
| Mid-June - early July*    | Paramyxo-herpes <b>BOOSTER (HERPES)</b>     |



### Need advice?

Your Neorni veterinarian will be happy to help

**TIMING** The sequence and repetition of vaccines are partly determined by the infection pressure in your loft. Contact NeorniVet for guidance on this.

**YOUNG PIGEON DISEASE** A **double vaccination** against **rotavirus**, given 3-4 weeks apart, provides the best protection against young pigeon disease (commonly referred to as 'adeno'). When infection pressure is high, a **third booster** may be given 3 weeks before training. Vaccinations against adenovirus and circovirus are not recommended because their effectiveness is unproven and there is a lack of pigeon strains.

**SUPPORT** Support your youngsters during the vaccination period in the first 6 weeks after weaning with **Immuno-Plus**, **Dysbac Liquid** and **Trichoryza Drops**.



## Old pigeons and yearlings

|               |                                                                          |
|---------------|--------------------------------------------------------------------------|
| Mid-February  | Smallpox (injection/follicle method) <b>CERTAINLY BENEFITS YEARLINGS</b> |
| Late February | <b>Paramyxo-herpes</b>                                                   |
| 14 days later | <b>Salmonella/paratyphoid**</b>                                          |

**SUPPORT** Get your old pigeons and yearlings off to the best start of the new season with **Trichoryza Drops**, **Dysbac Liquid** and **Ready**.



## Breeding pigeons:

|                           |                                 |
|---------------------------|---------------------------------|
| 6-7 weeks before breeding | <b>Salmonella/paratyphoid**</b> |
| Mid-October               | <b>Paramyxo-herpes</b>          |

**SALMONELLA/PARATYPHOID** Depending on infection pressure and immunity, it may be necessary to treat pigeons before vaccination, and vaccinate once or twice.

**HERPES** In case of herpes problems during the rearing of young pigeons, it is recommended to vaccinate breeding hens twice against herpes.

**PARAMYXO** An annual paramyxo vaccination is mandatory for breeding pigeons.

**SUPPORT** Stimulate breeding pigeons 2 weeks before pairing until the first clutch with **Vita-Min-DCP** and **Immuno-Plus**. Poor breeders/old cocks: supplement with **Vit E + Se**.



\* 3-4 weeks before the critical period

\*\* In case of previous infection, possibly after preventive treatment





# SCHEDULES FOR PIGEONS



# Breeding



## 2 weeks prior to mating until laying

| Trichoryza Drops                            | Vita-Min-DCP                                          | Immuno-Plus                                       |
|---------------------------------------------|-------------------------------------------------------|---------------------------------------------------|
| 10 ml/kg feed                               | 2 ml/l                                                | 2 ml/l                                            |
| During brooding                             | 2 weeks prior to mating until the first clutch        | 2 weeks prior to mating until the first clutch    |
| For 3 consecutive days                      | 2 days/week                                           | Daily                                             |
| Proven efficacy in resistant trichomoniasis | Extra calcium, phosphorus, Vit D3, and trace minerals | Stimulation of sperm cell and egg cell production |

## During brooding

56

| Tricho Geel                                                          |
|----------------------------------------------------------------------|
| 3 g/l or 6 g/kg feed                                                 |
| During brooding                                                      |
| For 5 consecutive days                                               |
| Reduce pressure of trichomoniasis + avoid hexamitiasis in youngsters |



## From hatching to weaning

| Immuno Booster NEW+ Feather Energy | Trichoryza Drops                            | Aminovital                                           |
|------------------------------------|---------------------------------------------|------------------------------------------------------|
| 20 g/kg feed + 5 ml/kg feed        | 10 ml/kg feed                               | 2-5 ml/l                                             |
| From hatching to weaning           | Full breeding period                        | From hatching to weaning                             |
| 2-3 days/week                      | For 1 day every 2 weeks                     | 3 days/week                                          |
| All-in premix                      | Proven efficacy in resistant trichomoniasis | Additional amino acids and vitamins for rapid growth |



# Flight preparation

## OLD PIGEONS AND YEARLINGS



**Trichoryza Drops**



**Dysbac Liquid**



**Ready**



10 ml/kg



1 day/2 weeks



Proven efficacy in resistant trichomoniasis



Insufficient training: **Immuno-Plus (2 ml/l) + Ready (2 ml/l)** for 14 days

2 ml/l

2 days/week

Crop, proventriculus and intestinal health, solid faeces

5 ml/l

Daily for 2 weeks before the start of the racing season

Form and conditioning

## YOUNG PIGEONS 'JUNIOR'

57



**Immuno Booster NEW**



**Trichoryza Drops**



**Dysbac Liquid**



20 g/kg



2-3 days/week for 6-8 weeks, afterwards 1 day/week throughout the flying season



Growth and energy, prevents nutritional deficiencies



Insufficient training: **Immuno-Plus (2 ml/l) + Ready (2 ml/l)** for 14 days

10 ml/kg

1 day/2 weeks

Proven efficacy in resistant trichomoniasis

2 ml/l

2 days/week

Crop, proventriculus and intestinal health, solid faeces



### Need advice?

Your Neorni veterinarian will be happy to help

►►► This schedule is an indication of how the NeorniPharma products can be used in certain circumstances. We try to develop a complete and appropriate schedule with as few products as possible. All these supplements can be used in combination, if required. In case of poor performance, we will look for the cause with you and decide upon any adjustments or further treatments. **Attention:** food supplements are best prepared the day before so they can be well absorbed, which will help the pigeons digest the food better.





# Flying season

## LOW SPEED



Trichoryza Drops



Dysbac Liquid



10 ml/kg

2 ml/l



Day after return home

Start of the week



For 1 day

For 2 days



Proven efficacy in resistant trichomoniasis

Crop, proventriculus and intestinal health, solid faeces



Pinch nostril, add drop, allow bird to inhale

Can be administered daily

58

## HIGH SPEED - MEDIUM DISTANCE



Aminovital



Trichoryza Drops



Dysbac Liquid



Ready



5 ml/l

10 ml/kg

2 ml/l

5 ml/l



Return home

Day after return home

Start of the week

Before basketing



For 1 day  
Demanding flights: for  
2-3 days

For 1-2 days

For 2 days

Great speed: 1 day  
Medium distance: 2-3 days



Fast recovery

Proven efficacy  
in resistant  
trichomoniasis

Crop, proventriculus  
and intestinal health,  
solid faeces

Form



In hot weather:  
basketing day

Pinch nostril, add drop,  
allow bird to inhale

Can be administered  
daily

Before an important  
race: daily for a period of  
10 days in combination  
with Immuno-Plus





# LONG DISTANCE

## Recovery



**Immuno Recup Caps**



**Aminovital**



**Immuno Booster NEW**



1 capsule/pigeon

5 ml/l

20 g/kg



For 1-2 days

For 2-3 days

For 2 days



Fast recovery

Fast recovery

Quick recovery, replenishing nutritional deficiencies

## Maintenance



**Trichoryza Drops**



**Dysbac Liquid**



**Respicor**



10 ml/kg

2 ml/l

3 ml/l



For 2 days

For 2 days

For 2 days



Proven efficacy in resistant trichomoniasis

Crop, proventriculus and intestinal health, solid faeces

Respiratory tract, white noses

59

## Boost before basketing



**Feather Energy**

**Vita Energy**

**Immuno-Plus**

**Red Race**



5 ml/kg

10 g/kg feed

2 ml/l

5 g/2 l



For 2-5 days

For 2-5 days

For 2-5 days

For 2-5 days



High-fat energy

Improves condition

Extra energy

Additional nutrient uptake



# Moulting Support



## Aminovital

2 ml/l

3 days/week

NO ACIDS + VITAMINS



## Immuno Booster NEW

20 g/kg

2 days/week

ALL-IN PREMIX



## Feather Energy

5 ml/kg

2 days/week

FATTY ACIDS



## Treating ectoparasites

### V-Exmite Drops

2 drops on the neck

Moulting period

DIVIDUAL TREATMENT



### Vermi-Complete

1 ml/l bath water

Moulting period

GROUP TREATMENT



60

## Preparation for breeding

Once the pigeon season finishes in late September/early October, we strongly recommend you **collect mixed faeces from the breeders for 5 days and have it analysed**. This mixed faeces is checked for worms, coccidiosis, and yeasts/fungi, following which a culture is grown to check for paratyphoid/salmonella. The results can be used to draw up a vaccination plan.

|                            | Paratyphoid/salmonella NEGATIVE | Paratyphoid/salmonella POSITIVE               |
|----------------------------|---------------------------------|-----------------------------------------------|
| <b>Treatment</b>           | Not necessary                   | Required for 10 days according to antibiogram |
| <b>Vaccination</b>         | Strongly recommended            | Required                                      |
| <b>Time of vaccination</b> | 6-8 weeks before pairing        | 2 days after treatment                        |



### Need advice?

Your Neorni veterinarian will be happy to help

**Discuss a plan of action tailored to your loft or problem with our veterinary practice NeorniVet (+32 (0)11 96 91 02)**

### Faecal analysis

- Collect faeces every day for 5 days and refrigerate it.
- Send the faeces, or take your sample to your veterinarian
- If required, the pigeons can be checked at the same time
- This is not harmful to moulting pigeons





# SCHEDULES FOR PSITTACIDAE



# Psittacidae

## PARROTS AND PARAKEETS

### General health



To prevent  
chronic  
deficiencies

Throughout the year  
1-2 days/week  
**Vita-Plus** or **Vita-Energy**

### Breeding

#### PREVENTION

6-8 weeks  
prior to  
breeding

Control mixed faeces from 3-5 days (vet)

- If necessary, deworm with **Vermi-Complete**
  - Group: 5 ml/l drinking water
  - Individual: 1 ml/kg body weight in the beak

Control ectoparasites

- Group: **Vermi-Complete** 1 ml/l bath water
- Individual:
  - <250 g: **Endo-Ecto Kanarie**
  - > 250 g: **V-Exmite Drops**

#### STIMULATES

daily 4-6  
weeks before  
breeding

**Vita Energy + Immuno-Plus**  
With drinking water and/or  
feed



#### REARING

daily from  
hatching to  
autonomy

Natural brooding:

- **Immuno Booster NEW** 2 g/100 g egg food/feed + **Immuno-Plus** 0.2 ml/100 ml drinking water

Hand rearing:

- **Immuno Booster NEW** 2 g/100 g soft food once per day + **Immuno-Plus** 1 ml/kg body weight once per day



# Crop and digestive problems



**For 7-14 days**  
1 ml/kg body weight  
**Hepa-Plus**



**For 7 days:**  
1 capsule/kg body weight  
**SBS Caps**



**Need advice?**

Your Neorni  
veterinarian will be  
happy to help

63

## Veterinarian check-up

- Disease testing through blood - feathers - cloacal swab
- Crop swab

# Moulting - Feathers



Additional  
support for  
spring growth

**Moulting + feather loss**  
3 days/week  
**Aminovital + Condi-Plus**  
2 ml/l 2 ml/l

# SilverSpray

**New** disinfectant spray

Bird-friendly  
Broad-spectrum  
Ready-made  
Multi-purpose





# SCHEDULES FOR CHICKENS

# Vaccination schedule



## Chicks

### Support

As required:

- First 14 days of life **Immuno Booster NEW** (20 g/kg feed)
- General vitamins: 3 days/week **Vita-Plus** (2 ml/l)
- Intestinal health + prebiotics: 2 days/week **Dysbac Liquid** (2 ml/l)
- Extra growth + amino acids: 2 days/week **Aminovital** (2 ml/l) + **Immuno-Plus** (2 ml/l)

### Vaccinations

**GOLD STANDARD:** 3 vaccines (Marek, NCD and smallpox)

|                       |                                           |
|-----------------------|-------------------------------------------|
| D1 (1st week of life) | Marek (2 strains) + IBD + NCD (injection) |
| D1 (1st week of life) | Infectious bronchitis (eye drop)          |
| 1st week of life      | Coccidiosis (drinking water)              |



**Need advice?**  
Your Neorni veterinarian will be happy to help

## Growing chickens (from 7 weeks) and full-grown chickens

### 66 Support

- General vitamins: **Vita-Plus** (2 ml/l) Prevention: 2 days/week After treatment: for 7 days.
- Intestinal health + prebiotics: **Dysbac Liquid** (2 ml/l). Prevention: 2 days/week In the case of flatter faeces: for 7 days.
- Respiratory tract support: **Respikor** (3 ml/l). Prevention: 2 days/week In the case of respiratory problems: for 7-14 days.

### Vaccinations

|                    |                                                              |
|--------------------|--------------------------------------------------------------|
| May-June (annual!) | Smallpox (+ ILT)                                             |
| May-June (annual!) | NCD (injection/drinking water)                               |
| August-September   | NCD (injection oil) <b>Mandatory for exhibition and sale</b> |
| August-September   | Infectious bronchitis (drinking water)                       |

## Advice for birds of all ages

- **GOLD STANDARD:** 3 vaccines (Marek, NCD and smallpox). Other vaccines are relevant to specific problem areas, and not routinely recommended.







# Advice and tips

## General health



**Throughout the year**  
1 day/week  
**Vita-Plus (2 ml/l)**

**During moulting**  
3 days/week  
**Aminovital and Condi-Plus (2 ml/l)**

## Weak egg shells

**Curative**

**Vita-Min-DCP** for 21 days

- Group: 5 ml/l or 10 ml/kg feed
- Individually through the beak: 1 ml/kg body weight

Worming day 1, day 7 and day 21

- Group: **Vermi-Complete** (5 ml/l)
- Individually through the beak: **Endo-Ecto Plus Caps**: 1 capsule/kg body weight, or **Vermi-Complete** 1 ml/kg body weight

**Prevention:**

Check mixed faeces every 3-4 months:

- Make sure you also have mixed faeces from 2-3 days analysed for endoparasites - these may be responsible for deficiencies, and cause irregularities in eggshells

Vaccination RT+IBmulti+ND+EDS

- Viral infections and malformations near the fallopian tube



67

## Snot



**Mild infection + new animal**  
7-14 days  
**Respisor (3 ml/l)**

## Wry neck/stargazing



**Wry neck**  
14-21 days  
**Vit E + Se**

**Group:**  
2 ml/l  
**Individual (beak):**  
0.2-0.5 ml/kg body weight

## Exhibition or show



**General form**  
7 days before the show  
**Immuno-Plus (2 ml/l)**

**Yellower legs**  
14-21 days before the show  
**PURE Yellow 1-2 g/kg**

If problems persist, make an appointment with your vet

# Export prices

Please find the NeorniPharma export prices for private individuals and professionals below. These prices are **exclusive of VAT**. The VAT rates for the country of destination are applied when shipping. As always, professionals are VAT exempt.

| PRODUCT                         | €<br>(excl.<br>VAT) | PRODUCT                           | €<br>(excl.<br>VAT) |
|---------------------------------|---------------------|-----------------------------------|---------------------|
| Immuno Booster NEW (1 kg)       | 53.72               | Trichoryza Drops (50 ml)          | 14.05               |
| Immuno-Plus (100 ml)            | 20.66               | Trichoryza Drops (250 ml)         | 48.76               |
| Immuno-Plus (250 ml)            | 45.45               | Trichoryza Drops (1 l)            | 136.36              |
| Immuno-Plus (1 l)               | 147.93              | Aminovital (100 ml)               | 15.70               |
| Immuno Recup Caps (80 capsules) | 19.83               | Aminovital (250 ml)               | 28.93               |
| Super Start-Up (200 g)          | 20.66               | Aminovital (1 l)                  | 53.72               |
| Vit E + Se (100 ml)             | 14.05               | Condi-Plus (250 ml)               | 21.49               |
| Vit E + Se (250 ml)             | 28.93               | Vita Energy (400 g)               | 24.79               |
| Hepa-Plus (100 ml)              | 16.53               | Vita-Min-DCP (100 ml)             | 16.53               |
| Hepa-Plus (250 ml)              | 29.75               | Vita-Min-DCP (250 ml)             | 28.93               |
| Hepa-Plus (1 l)                 | 65.29               | Vita-Min-DCP (1 l)                | 57.85               |
| Ready (100 ml)                  | 18.18               | Vita-Plus (100 ml)                | 14.88               |
| Ready (250 ml)                  | 39.67               | Vita-Plus (250 ml)                | 23.14               |
| Dysbac Liquid (250 ml)          | 30.58               | Vita-Plus (1 l)                   | 53.72               |
| Feather Energy (400 ml)         | 15.70               |                                   |                     |
| PikStop (100 ml)                | 23.14               | PURE                              |                     |
| Respikor (100 ml)               | 15.70               | Brewer's yeast (1 kg)             | 14.88               |
| Respikor (250 ml)               | 28.93               | Dextrose (1 kg)                   | 14.88               |
| Neorni Stabilizer (1 l)         | 16.53               | Electrolyte (500 g)               | 14.88               |
| Probi-Plus (150 g)              | 14.88               | Yellow (100 g)                    | 12.40               |
| SilverSpray (500 ml)            | 12.40               | Yellow (500 g)                    | 49.59               |
| Skin Spray (200 ml)             | 14.05               | Highly digestible protein (400 g) | 20.66               |
| CU-For Liquid (250 ml)          | 12.40               | Red (100 g)                       | 23.14               |
| SilverFlox (1 l)                | 26.45               | Red (500 g)                       | 70.25               |

# Index

|                                   |       |
|-----------------------------------|-------|
| Aminovital .....                  | 38    |
| Brewer's yeast .....              | 44    |
| Chicken schedules .....           | 66-67 |
| Condi-Plus.....                   | 39    |
| CU-For Liquid .....               | 34    |
| Dextrose .....                    | 44    |
| Dysbac Liquid .....               | 19    |
| Dysbacteriose .....               | 18    |
| Electrolyte .....                 | 44    |
| Feather Energy .....              | 22    |
| Fertility therapy .....           | 12    |
| Hepa-Plus .....                   | 14    |
| Highly digestible protein .....   | 44    |
| Immuno Booster NEW .....          | 7     |
| Immuno-Plus .....                 | 8     |
| Immuno Recup Caps .....           | 9     |
| Natural breeding promotion .....  | 6     |
| Neorni Stabilizer .....           | 28    |
| Pigeon schedules .....            | 56-60 |
| Pigeon vaccination schedule ..... | 54    |
| PikStop .....                     | 24    |
| Probi-Plus .....                  | 30    |
| Psittacidae schedules .....       | 62-63 |
| Ready .....                       | 15    |
| Red .....                         | 44    |
| Respikor .....                    | 25    |
| SilverFlox .....                  | 35    |
| SilverSpray .....                 | 31    |
| Skin Spray .....                  | 32    |
| Songbirds schedules .....         | 46-53 |
| Super Start-Up .....              | 10    |
| Trichoryza Drops .....            | 36    |
| Vita Energy .....                 | 40    |
| Vita-Min-DCP .....                | 41    |
| Vita-Plus .....                   | 42    |
| Vit E + Se .....                  | 11    |
| Yellow .....                      | 44    |

EN

T +32 (0)11 96 91 02  
E [info@neornipharma.be](mailto:info@neornipharma.be)  
W [www.neornipharma.be](http://www.neornipharma.be)

Front desk and depot:  
Beverlosesteenweg 129  
3583 Paal, Belgium

