



**2024**  
**NeorniPharma**  
supplements



# NeorniPharma Supplements

Preventive custom formulations for your birds

This catalogue is a comprehensive guide of the NeorniPharma range aimed at **disease prevention through natural, highly concentrated supplements**. This range offers a safe and successful alternative to medicinal treatments and a long-term solution. By targeting annually recurring problems, NeorniPharma aims to build immunity and consequently stronger birds.



## NeorniPharma Magistral Preparations

In addition to supplements, NeorniPharma also develops **Magistral preparations specifically aimed at disease control**. With this, NeorniPharma offers a short-term solution. Some of these products are featured in this catalogue. The full 'blue catalogue' is available from NeorniVet. Magistral preparations are only available on veterinary prescription.



NEW

### New front desk address

Beverlosesteenweg 129  
3583 Paal, Belgium

### New telephone number

+32 (0)11 96 91 02

### Additional dosages of powders

Discover alternative dosages on our 'Powders and capsules' dosage chart in this catalogue. View the full dosage chart at the back of the catalogue.



## Bird species in this catalogue



Pigeons



Songbirds



Fowl



Parrots



Birds of prey

Download our catalogues



All NeorniPharma products have unique formulas and are designed by veterinarians based on scientific research and practical experience.

## Breeding & immunity

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Immuno Booster NEW (1 kg)	20 g/kg (egg) feed	All-in premix	65
Immuno-Plus (100 ml)	2 ml/l	Breeding, show/competition, extra energy	25
Immuno-Plus (250 ml)	2 ml/l	Breeding, show/competition, extra energy	55
Immuno-Plus (1 l)	2 ml/l	Breeding, show/competition, extra energy	179
Immuno Recup Caps (80 c)	1 cap/0.5-1 KgBW	Immunity, recovery, energy	24
Super Start-Up (200 g)	5 g/kg feed	Start-up supplement for chicks, black spot	25
Vit E + Se (100 ml)	2 ml/l	Extra vitamin E & selenium	17
Vit E + Se (250 ml)	2 ml/l	Extra vitamin E & selenium	35

## Liver, muscles and condition

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Hepa-Plus (100 ml)	4 ml/l	Liver support	20
Hepa-Plus (250 ml)	4 ml/l	Liver support	36
Hepa-Plus (1 l)	4 ml/l	Liver support	96
Ready (100 ml)	5 ml/l	Conditioning preparation, fitness	22
Ready (250 ml)	5 ml/l	Conditioning preparation, fitness	48
Ready (1 l)	5 ml/l	Conditioning preparation, fitness	144

## Megabacteria and dysbacteriosis

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Dysbac Liquid (250 ml)	2 ml/l	Crop, proventriculus & intestinal health	37

## Natural oils

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Freeson Drops (30 ml)	1-2 drop/nostril	Unobstructed respiratory tract	17
Feather Energy (400 ml)	5 ml/kg feed	Flight Oil	29
PikStop (100 ml)	1 ml/l	Pecking	28
Respicor (100 ml)	3 ml/l	Respiratory tract, heat stress	19
Respicor (250 ml)	3 ml/l	Respiratory tract, heat stress	35

## Parasites

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Derma-Plus (50 ml)	8 ml/l/40 m <sup>2</sup>	Red mite, black lice	35
Exmite Spray (250 ml)	Spray	Ectoparasites	22
Exmite Spray (1 l)	Spray	Ectoparasites	69

## Probiotics and disinfection

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Neorni Stabilizer (1 l)	5-10 ml/100 ml	Probiotics – ambient	20
Probi-Plus (200 g)	3 g/l	Probiotics and prebiotics	18
SilverSpray (500 ml)	Spray	Disinfection spray	15
Skin Spray (200 ml)	Spray	Probiotics - skin infections	17

## Trichomonas, candida, aspergillosis & drinking water disinfection

PRODUCT	DOSAGE	INDICATION	PRICE (€)
CU-For Liquid (250 ml)	1-2 ml/l	Drinking water acidifier	15
SilverFloX (1 l)	2 ml/l	Drinking water disinfectant	32
Trichoryza Drops (50 ml)	1 drop/pigeon	Throat elixir	17
Trichoryza Drops (250 ml)	5 ml/l or 10 ml/kg	Throat elixir	59

## Vitamins, amino acids, electrolytes, minerals & trace elements

PRODUCT	DOSAGE	INDICATION	PRICE (€)
Aminovital (100 ml)	2 ml/l	Amino acids, electrolytes, vitamins	19
Aminovital (250 ml)	2 ml/l	Amino acids, electrolytes, vitamins	35
Aminovital (1 l)	2 ml/l	Amino acids, electrolytes, vitamins	85
Condi-Plus (100 ml)	2 ml/l	Moulting supplement	15
Condi-Plus (250 ml)	2 ml/l	Moulting supplement	26
Condi-Plus (1 l)	2 ml/l	Moulting supplement	83
Vita Energy (250g)	10g/kg feed	Conditioning Powder - vit, min and aa	25
Vita Energy (1kg)	10g/kg feed	Conditioning Powder - vit, min and aa	75
Vita-Min-DCP (100 ml)	2 ml/l	Vit D, calcium, phosphorus & other minerals	17
Vita-Min-DCP (250 ml)	2 ml/l	Vit D, calcium, phosphorus & other minerals	32
Vita-Min-DCP (1 l)	2 ml/l	Vit D, calcium, phosphorus & other minerals	88
Vita-Plus (100 ml)	2 ml/l	Water-soluble multivitamins	18
Vita-Plus (250 ml)	2 ml/l	Water-soluble multivitamins	28
Vita-Plus (1 l)	2 ml/l	Water-soluble multivitamins	83

## PURE

PRODUCT	PRICE (€)
Brewer's yeast (1 kg)	18
Dextrose (1 kg)	18
Electrolyte (500 g)	18
Yellow (100 g)	15
Yellow (500 g)	60
Highly digestible protein (400 g)	25
Red (100 g)	28
Red (500 g)	85

# NEORNI

NeorniPharma is part of Neorni, a team of veterinarians dedicated specifically to avian medicine. Neorni offers various services to ensure the best care for birds.

[www.neorni.be](http://www.neorni.be)  
[www.neornilab.be](http://www.neornilab.be)



NEORNIVET  
Veterinary practice



NEORNILAB  
Laboratory

- > Gender determination
- > Performance genes
- > Kinship analysis
- > Disease testing
- > Colour mutations



NEORNI PHARMA  
BY VETS FOR BIRDS  
Product range




Follow us  
on Facebook



Neorni (all services)  
Beverlosesteenweg 129  
3583 Paal, Belgium  
T +32 (0)11 96 91 02

---



# BREEDING & IMMUNITY



# Natural breeding boost

## Basic breeding

The basic breeding schedule is the essence of successful breeding. Several field trials showed that the birds were better prepared for breeding, which was reflected in more nests and more young per nest.



### Immuno-Plus

2 ml/l or 4 ml/kg

**Prior to breeding:**  
daily for a period of 6 weeks until 1st clutch

Breeding urge +++

Polypeptides

Stimulation of egg cell & sperm cell production



### Immuno Booster NEW

20 g/kg egg food/feed

**During breeding:**  
daily from hatching until weaning

Breeding urge +

Polypeptides & Premix  
breeding and rearing

Maintenance of egg cell & sperm cell production



### Need advice?

Your Neorni veterinarian will be happy to help



### Vit E + Se

2 ml/l or 4 ml/kg

Daily for 1-2w  
from breeding until the 1st clutch

Fertility +++

Linseed oil, arginine,  
spirulina, carnitine, zinc, etc

Extra high dose of  
vit E + Se

6

## Extra breeding (fertilisation)

Fertilisation problems occurring upon application of the basic schedules often have a nutritional or infectious cause.

▶▶ nutritionally speaking, a **Vit E + Se** supplement could certainly offer added value.

▶▶ In addition, certain germs with an important influence on fertilisation have been isolated in recent years. Manure culture before breeding can be carried out easily and quickly at NeorniVet veterinary practice(± 24h).

▶▶ In specific situations/species, extra added value can be achieved with the use of specific supplements such as **Vita-Plus, Ready, Hepa-Plus ...**

# IMMUNO BOOSTER NEW



## All-in premix for young birds



### Indication

- Highly concentrated premix
- Supports breeding results and maintains the condition of your birds
- Supply of extra energy
- Supports the immune system
- Helps build up intestinal flora
- Good development of the skeletal system
- Ideal for animals that are required to deliver top performances

### Administration 20 g/kg

- 2 large scoop (20g)/kg egg food/feed or 2 small scoop (2g)/100g egg food/feed
- During breeding season: add daily to egg food/feed
- Extra energy: 1-2 days per week

*Maize gluten, potato protein, fermented potato protein, yeast and its components, chicory root fibre, spirulina, yeast, trimagnesium citrate, mono-, di- and triglycerides from fatty acids (c4), tricalcium citrate, L-carnitine, betaine, Vit A, Vit D3, Vit E, Vit C, Vit B1, Vit B2, calcium D-pantothenate, Vit B6, Vit B12, Vit K, niacinamide, folic acid, choline, clinoptilolite, copper, iron, zinc, manganese, potassium iodide, selenium, cobalt, lysine, methionine, glycine, echinacea purpurea, turmeric, enterococcus faecium*

7

2 / 100 g





## Breeding, show/competition, extra energy



100 ml €25  
250 ml €55  
1 l €179

8

### Indication

- Supports breeding results and maintains the condition of your birds
- Supply of extra energy
- Good appetite and feather patterns
- To be used before/after intense exertion/competition
- Supports the development & growth of young animals

### Administration 2 ml/l drinking water

- Breeding: Daily, starting 6 weeks before breeding Stop supplementation after the 1st clutch After hatching, resume daily supplementation to the parents (and the young) until leaving the nest
- Show/competition: starting 5 days before show/competition, daily 2 ml/l drinking water or 2 ml/kg body weight directly into the bill
- Supply of extra energy: 2 ml/l drinking water for a period of 14 days or 2 ml/kg body weight directly into the bill until recuperated

# IMMUNO RECUP CAPS



## Immunity, recovery, energy



80 capsules

€24

### Indication

- Stimulates recovery after illness/exertion
- Supply of extra energy
- Supports the immune system
- Ideal for top performance

### Administration

- Recovery/immunity boost: 1 capsule/0.5-1kg orally for 5-10 days
- Extra energy: 1 capsule/0.5-1kg orally
- Pigeon: day of return home, 1 capsule/pigeon
- To support breeding: for a period of 14 days before mating until laying 1 capsule/0.5-1kg
- *Immuno Recup Caps* can be administered directly or with the food. When administered via food, the full dose should be consumed.

*Maize gluten, potato protein, fermented potato protein, yeast and its components, chicory root fibre, spirulina, yeast, trimagnesium citrate, mono-, di- and triglycerides from fatty acids (c4), tricalcium citrate, L-carnitine, Vit A, Vit D3, Vit E, Vit C, Vit B1, Vit B2, calcium D-pantothenate, Vit B6, Vit B12, Vit K, niacinamide, folic acid, choline, clinoptilolite, copper, iron, zinc, manganese, potassium iodide, selenium, cobalt, lysine, methionine, glycine, echinacea purpurea, turmeric, enterococcus faecium*



## Start-up supplement for chicks, black spot



200 g €25

10

### Indication

- Smooth start for your young chicks in the first 6 weeks of life
- Natural suppression of bacterial and necrotic enteritis/diarrhoea
- Rapid growth and larger birds due to smooth start and natural suppression of black spot

### Administration 5 g/kg egg food/feed

- Daily throughout the first 6 weeks of life

# VIT E + SE



## Extra vitamin E + selenium



100 ml €17

250 ml €35

+ spirulina  
+ zinc  
+ linseed oil

### Indication

- For animals with a deficiency of vitamin E selenium
- Recommended for nervous symptoms

### Administration

- 2 ml/l drinking water or 1 ml/kg body weight directly into the bill
- Recommended for nervous symptoms: 7-10 days




#### Fertility +++

2 ml/l or 4 ml/kg feed daily for 1-2 weeks,  
before breeding until first clutch

*Vit E (synthetic + natural and more efficiently absorbed form), selenium, linseed oil, spirulina, carnitine, arginine, zinc glycinate*





# LIVER, MUSCLES & FITNESS





## Liver support



100 ml €20  
250 ml €36  
1 | €96

14

### Indication

- Supports liver function
- Relieves stress on the liver

### Administration

- Curative: 4ml/l drinking water for a period of 5 days
- Prevention: 2 ml/l drinking water, 2 to 3 days per week



Can be administered daily to support the liver during administration of medication or dye



## Conditioning preparation, fitness



100 ml €22  
250 ml €48  
1 l €144

### Indication


- To be used when fitness decreases or when improving the overall condition
- Extra source of iron, iodine, L-carnitine and vitamin B12
- Stimulates fat metabolism and endurance
- For the improvement of flight performance
- Supports blood circulation

### Administration 5 ml/l drinking water

- In preparation for an important competition: daily for a period of 2 weeks
- Overweight birds: 3-5 days per week until desired results are achieved
- In support of anaemia: 7-10 days
- Avoid administration at temperatures above 30°C

*Glycerol, L-carnitine, red beet juice, potassium iodide, ferrous glycinat, Vit B12, magnesium citrate, sodium chloride, magnesium chloride, calcium chloride, glucose, vitamin E, selenium, zinc, copper, manganese, vitamin A, vitamin B1, vitamin B2, vitamin B5, vitamin B6, Vitamin PP (niacinamide), vitamin D3, vitamin C, mix of natural aromatic components*





# MEGABACTERIA & DYSBACTERIOSIS



# Dysbacteriosis

crop

proventriculus

intestines



**Dysbac Liquid**

prevention of  
dysbacteriosis

**Dysbac Liquid** is an improved & extended version of **CU-For Liquid** with more focus on **intestinal health** and **general immunity**

18



2 ml/l
+
+++
+++
+++
+++
+++
+++
CROP, PROVENTRICULUS & INTESTINAL HEALTH, PREVENTION OF DYSBACTERIOSIS

- ▶▶ **Short-chain fatty acids:** formic acid, propionic acid, citric acid
- ▶▶ **Medium-chain fatty acids:** butyric acid, mono-, di- and triglycerides of fatty acids (C8-C10)
- ▶▶ **Aromatic substances:** oregano, echinaceae purpureae, ginger, cinnamon, black pepper, garlic, and boswellia
- ▶▶ **Vitamin C**
- ▶▶ **Chelated copper and zinc**
- ▶▶ **Extra fibre**
- ▶▶ **Fish oil**

- DOSAGE
- ACIDIFICATION OF DRINKING WATER (pH)
- CROP HEALTH
- POVENTRICULUS HEALTH
- INTESTINAL HEALTH
- GENERAL IMMUNITY

PURPOSE



1-2 ml/l
+++
+++
++
-
-
WATER ACIDIFICATION, WATER QUALITY

- ▶▶ **Short-chain fatty acids:** ammonium formate, acetic acid, lactic acid, formic acid, propionic acid
- ▶▶ **Copper and zinc sulphates**



**Need advice?**

Your Neorni veterinarian will be happy to help



## Crop, proventriculus and intestinal health



250 ml €37

### Indication

- Forms a protective film over the intestinal mucosa
- Increases the animal's immunity
- Helps to maintain the natural balance of intestinal flora
- To be used in cases of watery droppings

### Administration 2 ml/l drinking water

- Prevention: 1 to 2 days per week
- In case of digestion problems: for a period of 7-14 days



### Prevention of megabacteriosis & dysbacteriosis

- 1-2 days a week 2 ml/l drinking water
- May be combined with ACS

*Monobutyryn, prebiotic fibre, fish oil, medium-chain fatty acids, sodium chloride, glycerol, zinc, copper, formic acid, propionic acid, citric acid, 10 different strains of oregano, echinacea purpurea, cinnamon, thyme, Boswellia, black pepper, microfused essential oils, vitamin C*



---



# NATURAL OILS



## Clear respiratory tract



22

### Indication

- Hardened snot and black nares

### Administration 1-2 drops per nostril

- All birds: for respiratory problems: 2-3 times/day for 7 days.
- Preventive in pigeons: 14 days before the flying season: for 3 days. The day before basketing. The week before an important flight: daily for 3 days.



## Flight oil, source of energy from lipids



### Indication

- Source of energy from lipids
- Feather quality, optimal feathering
- Extra vitamins A, D, and E
- Hoge source of omega 3-6-9 fatty acids

### Administration 5-10 ml/kg feed

- During moulting: 2 days per week
- Before and during breeding: 2-3 days per week
- During racing season: 1-2 days per week
- *Feather Energy* can be combined with *Vita Energy* or *Immuno Booster NEW*



Can be used to improve the absorption of *Immuno Booster NEW* and *Vita Energy* with the feed

*Cold-pressed oils: vegetable oils and fats (linseed oil), cod liver oil, rapeseed oil, salmon oil, crude lecithins, corn germ oil, wheat germ oil, sunflower oil, hemp seed oil (including Vit A, D, E)*



## Pecking, unrest



100 ml €28

NEW

24

### Indication

- Pecking in birds and poultry

### Administration

- 1 ml/l drinking water for 2-4 weeks



#### Feather picking/pecking

Serve simultaneously with *Immuno-Plus* for optimal results



## Respiratory tract, heat stress



100 ml €19

250 ml €35

NEW  
formerly  
Herbi+E

### Indication

- Natural solution to support respiratory problems
- Suppression of pain/inflammation in case of respiratory problems
- Stuck slime
- Heat stress

**Administration** 3 ml/l drinking water or 0.5 ml/kg body weight directly into the bill

- Preventive: when purchasing new animals: for 10-14 days
- Curative: for mild to moderate sniffing of chickens: for 10-14 days
- Heat stress: 3-5 ml/l
- Additional support: *Respicor* can be combined with medication for a soothing effect
- If problems persist: contact your vet

# NEW

# neofalco

products for birds of prey



more info: [info@neornipharma.be](mailto:info@neornipharma.be)



# PARASITES





## Red mite, black lice



50 ml €35

28

### Indication

- Spray for surroundings (coop) and birds to combat haematophagous mites
- Not harmful to birds: spray may be applied directly to birds; coop may be treated in the presence of the birds

### Administration 8 ml/l/40 m<sup>2</sup>

- Spray: 8 ml/l water for 40 m<sup>2</sup> coop
- Repeat after 1 week

# EXMITE SPRAY



## Ectoparasites



250 ml €22

1 | €69

### Indication

- Ectoparasites

### Administration

- Red mite/feather lice/black lice: Spray at the level of the neck, tailbone and cloaca, and under the wings. Once, repeat after 7 days.
- Scaly leg mites: Spray the area to be stabilized daily for 14 days.
- Feather mite: Spray at the level of the neck, tailbone and cloaca, and under the wings. Once, repeat after 10 days and after 20 days.
- Preparation for breeding: Preventively spray nests/ breeding boxes/coops/perches with *Exmite Spray*. May be repeated weekly in case of high load.




#### Heavy feather mite infections

Day 0, 10 and 20: serve simultaneously *V-Exmite Drops* and/or *Endo-Ecto Caps* for best results



---



# PROBIOTICS & DISINFECTION

# SilverSpray

**New** disinfectant spray

Bird-friendly  
Broad-spectrum  
Ready-to-use  
Multi-purpose



# NEORNI STABILIZER



## Probiotic ambient spray



### Indication

- Reduction of pathogenic load in your birds' surroundings
- Forms a stable and healthy environment by occupying it with safe, probiotic bacteria

### Administration

- To be used diluted: 5-10% (5-10 ml/100ml) in lukewarm water
- Mist the entire area to be stabilized
- First week, spray the coop daily (may be done in the presence of the birds, and even the nestlings).
- Subsequently, spray twice a week to maintain probiotic protection
- The product diluted with water has a maximum shelf life of 5 days.



## Probiotics and prebiotics



34

### Indication

- Increases the beneficial flora in the crop and the intestines
- Ensures improved digestion and supports the initiation of the crop and intestinal flora in chicks
- After a course of antibiotics

### Administration 3g/L drinking water or 6g/kg feed

- Preventive: for 2 days/week
- Wet droppings or after antibiotic treatment: for 5-7 days
- Refresh daily
- May be combined with other supplements (vitamins)





## Disinfectant spray



### Indication

- Broad-spectrum disinfectant
- Bactericide - Fungicide - Virucide

### Administration

- Ready-to-use disinfectant spray
- Suitable for pens, cages, equipment and eggs



Safe to apply in the presence of your birds



## Probiotic spray for skin infections



200 ml €17

36

### Indication

- Suitable for birds with sensitive skin
- To be used externally for care of the skin
- Reduces the formation of scaly skin


### Administration

- Mist the area to be stabilized



#### **Powder comb/flavus/white flakes**

Spray the comb or infected area daily. Not harmful to eyes and respiratory tract.



# TRICHOMONIASIS, CANDIDA ASPERGILLOSIS & DRINKING WATER DISINFECTION





## Acidification of drinking water



250 ml €15

38

### Indication

- Acidification of drinking water, lowering pH

### Administration 1-2 ml/l drinking water

- Prevention: 1 to 2 days per week



Exceptionally concentrated formula, very economical to use



## Drinking water disinfection



### Indication

- Water disinfection
- Prevention of bad bacteria, viruses, yeasts and moulds
- Immunity support

### Administration

- 2ml/L drinking water



Longer and better action due to silver addition



## Throat elixir



50 ml €17  
250 ml €59

40

### Indication

- Preventive approach to trichomoniasis
- Supports throat and crop health
- Positive effect on the upper respiratory tract
- Plays a role in resistance to disease

### Administration

- The morning after returning home:
  - 1 drop in the throat and/or 1 drop per naris
  - 10ml/kg feed or 5ml/l drinking water
- Prevention: 1 day per week.
- At high pressure: 2 days per week
- Caution: stains clothing and hands



Proven efficacy in resistant trichomonas cases:  
for 5 days 5-8 ml/l drinking water

---



# VITAMINS, AMINO ACIDS, ELECTROLYTES, MINERALS & TRACE MINERALS





## Amino acids, electrolytes & vitamins



100 ml €19  
250 ml €35  
1 | €85

Extra vitamin E,  
selenium & BCAA  
amino acids

42

### Indication

- All essential amino acids crucial for proper moulting/feather development/pigmentation
- Stimulates recovery/repair
- Limits muscle loss

### Administration 2 ml/l drinking water

- Racing pigeons: 5 ml/l during the racing season
- Throughout the year: 1 to 2 days per week
- Throughout the entire moulting period: 3 days/week. May be combined with *Condi-Plus*.



**Additional support for racing pigeons in warm weather**

5 ml/l drinking water, the day of basketing

*Dextrose, glycerol, sodium chloride, probiotic fibre, trimagnesium citrate, red beet juice, potassium chloride, saccharose, calcium chloride, tricalcium citrate, lysine, methionine, threonine, tryptophan, histidine, carnitine, betaine, choline, Vit B1, Vit B2, niacinamide, calcium D-pantothenate, Vit B6, Vit B12, folic acid, biotin, Vit C, Vit E, selenium, copper, zinc, manganese, mix of natural aromatic components*



## Moulting supplement



100 ml €15  
250 ml €26  
1 l €83

### Indication

- Supports bird health during molting
- Supports the metabolism
- Supply of extra energy
- Increases appetite

### Administration 2 ml/l drinking water

- Throughout the entire molting period: 3 days/ week. May be combined with *Aminovital*.



## Conditioning powder – Vitamins, minerals and amino acids



44

### Indication

- Extra energy
- Extra vitamins, minerals and amino acids
- Increases endurance
- Demanding races, long-distance races

### Administration 10 g/l feed

- General: Throughout the year, 1-2 days/week. In case of poor/impaired condition: for a period of 14 days.
- Pigeon: Medium distance/long distance/in warm weather, 2-3 days before basketing. For weekly long-distance races/flights: daily from return home until basketing.



*Dextrose, maltodextrin, yeast and its components, chicory root powder, trimagnesium citrate, sodium bicarbonate, refined coconut oil, glucose syrup, caseinate, sodium chloride, potassium chloride, trisodium phosphate, tricalcium citrate, Vit A, Vit D3, Vit E, Vit K, Vit B1, Vit B2, Vit B3, Vit B5, Vit B6, Vit B11, Vit B12, biotin, Vit C, cobalt, iron, copper, zinc, manganese, potassium iodine, selenium, lysine, methionine, mix of natural aromatic components*

# VITA-MIN-DCP



## Liquid source of vitamin D, calcium, phosphorus and other minerals



100 ml €17  
250 ml €32  
1 l €88

Extra calcium  
& phosphorus

### Indication

- For animals with a deficiency in Vitamin D3, calcium, phosphorus and other minerals
- Supports bone formation and development in young animals
- Supports egg shell formation
- Poor egg shell quality/shell-less eggs

### Administration 2 ml/l drinking water

- Prevention: 2 ml/l
  - 2 weeks before breeding until the eggs are laid: 2-3 days per week
  - Throughout the first 6 weeks of life: 2-3 days per week
- Curative: 5 ml/l for a period of 7 days



Due to the highly concentrated formula, limited sedimentation may be visible in the drinking water. This does not affect the action.

*Trisodium phosphate, trimagnesium citrate, calcium chloride, magnesium chloride, tricalcium citrate, vitamin D3, cholecalciferol, zinc, manganese, copper, iron*



## Water-soluble multivitamins



46

### Indication

- Supplement containing all essential vitamins
- Overall daily health
- Prevention of chronic vitamin A deficiency

### Administration 2 ml/l drinking water

- Weekly: for 1 to 2 days per week
- After illness/recuperation: for a period of 3 to 5 consecutive days



For parrots and parakeets, we strongly recommend 1-2 days per week *Vita-Plus* via the drinking water 1-2 days a week to prevent vitamin A deficiency



High concentration of vitamin K helps prevent or repair bleeding




# SCHEDULES FOR SONGBIRDS





# Annual Euro & Canary schedule

## Timeline



### Preparation for breeding

- Breeding boost
- Liver support
- Black spot prevention

### Hatching to weaning

- Optimal start & sweat disease prevention
- Suppression of coccidiosis/atoplasmosis, bacteria, viruses, yeasts and fungi

### Weaned young

- Prevention coccidiosis/atoplasmosis, megabacteriosis and dysbacteriosis
- Limiting pressure with immunity build-up





# Annual Euro & Canary schedule

*The annual schedule is a general guide for preparation, breeding, hatching, weaning and moulting. Talk to your veterinarian about the specific requirements for your birds and devise a customised schedule together.*

## Preparation from 6-8 weeks prior to breeding

1. A healthy environment is the best start! (see CAS or SilverSpray)
2. Prevention of atoxoplasmosis/coccidiosis + megabacteriosis/dysbacteriosis
  - **Veterinarian:** bird check-up + manure tests 6-8 weeks prior to breeding
  - Only use courses of treatment in case of problems:
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 3 days Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
3. For a period of 6 weeks prior to breeding
  - **Immuno-plus (2 ml/l) daily**
    - Breeding boost; birds will be in top (breeding) condition
  - 1-2 days per week **Hepa-Plus (4 ml/l)**
    - Liver support
  - As required: Vita-Plus, Vita Energy, Ready, Vit E+Se, Vita-Min-DCP
    - These are suitable for use with Immuno-Plus
  - Extra: black spot prevention
    - For 7 days **Gould Mix (5 g/l)** during breeding preparation
4. Prevention of haematophagous mites
  - **Exzolt:** 1 ml/l for 1 day, repeat on days 7 and 14
    - Does not affect fertility & nestlings/young
    - Can be repeated every 3 months
  - Extra prevention of haematophagous mites
    - Before occupation, treat coop/loft with:
      - **Derma-Plus/Exmite Spray**
      - Can be repeated while the birds/nestlings/young are present

49

## Breeding: hatching to weaning

1. **Immuno Booster NEW (2 g/100 g or 20 g/kg egg food/feed)**
  - All-in premix for young birds
2. First 6 weeks
  - Option 1: **Super Start-Up (5 g/kg egg food/feed)**
    -  Natural suppression and prevention of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
  - Option 2: **Super Start-Up Blue (20 g/kg egg food/feed)**
    -  Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
3. Start from 2 days before hatching until 5 days after hatching **Coli-Mix (5 g/l)**
  - Prevention of mushy chick disease (Colibacillosis)
  - Alternative to Coli-Mix: **Gould Mix (5 g/l)**



## Breeding: weaned young

1. A healthy environment is the best start!

- **CAS or SilverSpray**
  - Environment free of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and moulds
- **Neorni Stabilizer:** Probiotic ambient spray
  - Prevention by reducing ambient pressure

2. Preventive cure

- 3rd week after weaning
  - **Increased susceptibility to atoxoplasmosis/megabacteriosis**
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 3 days Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)
    - 5 days Sulfaclozine (3 g/l) + Hepa-Plus (4 ml/l) + Megadysbac (5 g/l)

3. Then throughout the year (from weaning to breeding period)

- **ACS 3-5 days (0.5 ml/l) 1-2 a month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis

## 50 Moulting period

1. Throughout moulting **Condi-Plus + Aminovital:**

- Condi-Plus (2 ml/l): 3 days/week
- Aminovital (2 ml/l): 3 days/week
- May be administered simultaneously


2. Throughout the year (from weaning to breeding)

- **ACS 3-5 days (0.5 ml/l) 1-2 a month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis

At NeorniPharma, we always strive for the best long-term solution. We are constantly looking for **alternatives to anti-microbials** to limit the risk of resistance. The goal should be give our birds **as few preventive treatments with antimicrobial agents as possible**. Clinical research and any additional tests are crucial for this. After the diagnosis, an appropriate treatment can be administered. **Antibiotics are not harmless**. Consult your vet for advice about their correct and responsible use. **Autopsy:** don't simply throw away a bird that has died! The cadaver can provide a lot of information to ensure appropriate treatment can be sought. It is best to keep the bird cool (in a fridge, for example) and carry out an investigation within 24 hours (max. 72h).



# Preparation for breeding



From 6 weeks prior to breeding until the 1st clutch


Immuno-Plus	Hepa-Plus
2 ml/l	4 ml/l
6 weeks prior to mating until the 1st clutch	6 weeks prior to mating until the 1st clutch
Daily	1-2 days/week
Stimulation of sperm cell & egg cell production	Supports the liver



## Need advice?

Your Neorni veterinarian will be happy to help

51



Extra support

Vit E + Se	Vita-Min-DCP	Ready	Vita-Plus/Vita Energy
2 ml/l	2 ml/l	3-5 ml/l	2 ml/l or 10 g/kg
2 weeks prior to mating until the 1st clutch	2 weeks prior to mating until the 1st clutch	6 weeks prior to mating until the 1st clutch	Throughout the year
Daily	2 days/week	1-2 days/week	1-2 days/week
Extra-high doses of Vit E + Se, linseed oil, arginine, spirulina, carnitine, zinc, etc.	Extra calcium, phosphorus, vit D3, and trace minerals	Converts fat into usable energy for the bird	Extra vitamins

FERTILISATION

EGGSHELL

CONDITION

VITAMINS

▶▶▶▶ All products can be used in combination





# Breeding: hatching to weaning



52

**Coli-Mix** 

5 g/l or 10g/kg egg food/feed

---

2 days before to 5 days after hatching

---

Does not affect fertility

---

Broad-spectrum mix

---

**Prevention of mushy chick disease & Bengalese finch disease**

**Immuno Booster NEW**

20 g/kg egg food/feed

---

Daily from hatching until weaning

---

Positive effect on fertility

---

All-in premix

---

**Maintenance of egg cell & sperm production**

**Super Start-Up** 

5 g/kg egg food/feed

---

**Natural suppression and prevention of coccidiosis/ atoxoplasmosis, bacteria, viruses, yeasts and fungi**

**OR**

**Super Start-Up Blue**

20 g/kg egg food/feed 

---

**Suppression of coccidiosis/ atoxoplasmosis, bacteria, viruses, yeasts and fungi**

**Immuno Booster NEW** contains all essential components for **rearing young birds**. It is a unique **comprehensive** premix to mix under egg food/feed, providing all necessary nutrients in **exceptionally high concentrations**.

**Immuno Booster NEW Polypeptides** Keep the birds in form during breeding. **Saccharomyces cerevisiae** Source of protein, amino acids, and nucleotides. **Chicory root fibre** Prebiotic fibre: balanced digestion. **Spirulina** Source of protein, fats, carbohydrates, vitamins, and minerals. **Electrolytes** Fluid balance. **Fatty acids** Energy, balanced digestion. **All water-soluble and fat-soluble vitamins** General immunity, antioxidants, growth, energy release. **All amino acids** Building muscle and feather cover. **L-carnitine** Supports fat digestion and muscle building. **Choline** Liver support. **Minerals** (cobalt, iron, copper, zinc, manganese, iodine, selenium) General immunity, antioxidants, growth, energy release. **Clinoptolite** General health and intestinal health. **Probiotics** Intestinal flora stabilizer. **Turmeric and echinacea** General immunity and health.



# Breeding: weaned young

	COCCIDIOSIS & ATOXOPLASMOSIS	MEGABACTERIA & DYSBACTERIOSIS
	<b>ACS</b>	<b>Dysbac Liquid</b>
	0.5 ml/l	2 ml/l
	From weaning until next breeding time	From weaning until next breeding time
	3-5 days/month If required: 2x/month	1-2 days/week
	<b>COCCIDIOSIS &amp; ATOXOPLASMOSIS</b>	<b>MEGABACTERIOSIS, DYSBACTERIOSIS &amp; FLAGELLATES</b>



53

## Extra support throughout the year

SUPPORTS THE LIVER	GENERAL SUPPORT
<b>Hepa-Plus</b>	<b>Vita-Plus / Vita Energy</b>
2 ml/l	2 ml/l or 10 g/kg
Throughout the year	Throughout the year
1-2 days/week When adding dye: 2 ml/l daily	1-2 days/week
<b>LIVER SUPPORT</b>	<b>EXTRA VITAMINS</b>





# Annual Exotics schedule



*The annual schedule is a general guide for preparation, breeding, hatching, weaning and moulting. Talk to your veterinarian about the specific requirements for your birds and devise a customised schedule together.*

## Preparation from 6-8 weeks prior to breeding

1. A healthy environment is the best start! (see CAS or SilverSpray)
2. Prevention of flagellates (cochlosomosis) + campylobacteriosis
  - **Veterinarian:** bird check-up + manure tests 6-8 weeks prior to breeding
  - Only use courses of treatment in case of problems:
    - For 7 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)
3. For a period of 6 weeks prior to breeding
  - **Immuno-plus (2 ml/l) daily**
    - Breeding boost; birds will be in top (breeding) condition
  - 1-2 days per week **Hepa-Plus (4 ml/l)**
    - Liver support
  - As required: Vita-Plus, Vita Energy, Ready, Vit E+Se, Vita-Min-DCP
    - These are suitable for use with Immuno-Plus
  - Prevention of cochlosomosis + campylobacteriosis
    - For 7 days **Gould Mix (5 g/l)** during breeding preparation
4. Prevention of ectoparasites
  - Haematophagous mites
    - **Exzolt:** 1 ml/l for 1 day, repeat on days 7 and 14
      - Does not affect fertility & nestlings/young
      - Can be repeated every 3 months
    - Extra prevention of haematophagous mites
      - The day prior to occupation, treat coop/loft with:
        - **Derma-Plus/Exmite Spray**
          - Can be repeated while the birds/nestlings/young are present
  - Tracheal mites
    - **Vermi-Complete:** 2.5-5 ml/l for 1 day, repeat after 3 weeks

## Breeding: hatching to weaning

1. **Immuno Booster NEW (2 g/100 g or 20 g/kg egg food/feed)**
  - All-in premix for young birds
2. First 6 weeks
  - Option 1: **Super Start-Up (5 g/kg egg food/feed)**
    -  Natural suppression and prevention of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
  - Option 2: **Super Start-Up Blue (20 g/kg egg food/feed)**
    -  Suppression of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and fungi
3. Start from 2 days before hatching until 5 days after hatching **Coli-Mix (5 g/l)**
  - Prevention of mushy chick disease (Colibacillosis)
  - Alternative to Coli-Mix: **Gould Mix (5 g/l)**



#### 4. For problems in the first 6-8 weeks of life

- **Gould Mix (5 g/l)**
  - For the treatment of cochlosomosis (Bengalese finch disease), yeasts and bacterial diarrhoea

### Breeding: weaned young

#### 1. A healthy environment is the best start!

- **CAS or SilverSpray**
  - Environment free of coccidiosis/atoxoplasmosis, bacteria, viruses, yeasts and moulds
- **Neorni Stabilizer:** Probiotic ambient spray
  - Prevention by reducing ambient pressure

#### 2. Preventive cure

- 3rd week after weaning
  - **Increased susceptibility to coccidiosis/cochlosomosis/campylobacter**
    - 5 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)
    - 3 days: Hepa-Plus (4 ml/l)
    - 5 days Gould Mix (5 g/l) + Hepa-Plus (4 ml/l)

#### 3. Then throughout the year (from weaning to breeding period)

- **ACS 3-5 days (0.5 ml/l) 1-2 a month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis
  - Prevention of flagellates (cochlosomosis)

55

### Moulting period

#### 1. Throughout moulting **Condi-Plus + Aminovital:**

- Condi-Plus (2 ml/l): 3 days/week
- Aminovital (2 ml/l): 3 days/week
- May be administered simultaneously

#### 2. Throughout the year (from weaning to breeding)

- **ACS 3-5 days (0.5 ml/l) 1-2 a month**
  - Prevention of coccidiosis while improving immunity!
- **Dysbac Liquid 1-2 days per week (2 ml/l)**
  - Prevention of megabacteriosis & dysbacteriosis
  - Prevention of flagellates (cochlosomosis)

At NeorniPharma, we always strive for the best long-term solution. We are constantly looking for **alternatives to anti-microbials** to limit the risk of resistance. The goal should be give our birds **as few preventive treatments with antimicrobial agents as possible**. Clinical research and any additional tests are crucial for this. After the diagnosis, an appropriate treatment can be administered. **Antibiotics are not harmless**. Consult your vet for advice about their correct and responsible use. **Autopsy:** don't simply throw away a bird that has died! The cadaver can provide a lot of information to ensure appropriate treatment can be sought. It is best to keep the bird cool (in a fridge, for example) and carry out an investigation within 24 hours (max. 72h).





# SCHEDULES FOR PIGEONS





# Breeding



## 2 weeks prior to mating until laying

Vita-Min-DCP
2 ml/l
2 weeks prior to mating until the 1st clutch
2 days/week
Extra calcium, phosphorus, vit D3, and trace minerals

Immuno-Plus
2 ml/l
2 weeks prior to mating until the 1st clutch
Daily
Stimulation of sperm cell & egg cell production

**Need advice?**  
Your Neorni veterinarian will be happy to help

## During brooding

58

Trichoryza Drops
10ml/kg feed or 5ml/l
During brooding
For a period of 3 consecutive days
Proven effectiveness in resistant trichomonas



Fertilisation problems occurring upon application of the basic schedules often have a nutritional or infectious cause.

▶▶ Nutritionally speaking, a **Vit E + Se** supplement can certainly offer added value.

▶▶ In certain situations, extra added value can be created through the addition of certain supplements such as **Vita-MIN-DCP** (poor shell quality) and **Ready** (overweight pigeons).

## From hatching to weaning

Immuno Booster NEW+ Feather Energy
20 g/kg feed + 5 ml/kg feed
From hatching to weaning
2-3 days/week
All-in premix

Trichoryza Drops
10ml/kg feed or 5ml/l
Full breeding period
For 1 day every 2 weeks
Proven effectiveness in resistant trichomonas

Aminovital
2-5 ml/l
From hatching to weaning
3 days/week
Additional amino acids and vitamins for rapid growth



# Flight preparation

## Old pigeons and yearlings



**Trichoryza Drops**



**Dysbac Liquid**



**Ready**



5 ml/l or 10 ml/kg

2 ml/l

5 ml/l



1 day/2 weeks

2 days/week

Daily for 2 weeks before the start of the racing season



Proven efficacy in resistant trichomonas

Crop, proventriculus & intestinal health, solid droppings

Form & fitness building



Insufficient training: **Immuno-Plus (2 ml/l) + Ready (2 ml/l)** for a period of 14 days

## Young pigeons 'JUNIOR'



**Immuno Plus**



**Immuno Booster NEW**



**Trichoryza Drops**



**Dysbac Liquid**

59



2 ml/l

**OR**

20 g/kg

5 ml/l or 10 ml/kg

2 ml/l



Daily for 4 to 6 weeks

1 day/2 weeks

2 days/week



Growth & energy

Proven efficacy in resistant trichomonas

Crop, proventriculus & intestinal health, solid droppings



Insufficient training: **Immuno-Plus (2 ml/l) + Ready (2 ml/l)** for a period of 14 days



### Need advice?

Your Neorni veterinarian will be happy to help

>>> This schedule is an indication of how the NeorniPharma products can be used in certain circumstances. We try to develop a complete and appropriate schedule with as few products as possible. All these supplements can be used in combination, if required. In case of poor performance, we will look for the cause with you and decide upon any adjustments or further treatments.

**Attention:** food supplements are best prepared the day before so they can be well absorbed, which will help the pigeons digest the food better.



# Flying season

## Small speed



**Trichoryza Drops**



**Dysbac Liquid**



5 ml/l or 10 ml/kg or 1 drop/nostril

2 ml/l



Day after return home

Start of the week



For 1 day

For 2 days



Proven efficacy in resistant trichomonas

Crop, proventriculus & intestinal health, solid droppings



Pinch naris, add drop, allow bird to inhale

Can be administered daily

## Short distance - half distance

60



**Aminovital**



**Trichoryza Drops**



**Dysbac Liquid**



**Ready**



5 ml/l

5 ml/l or 10 ml/kg or 1 drop/naris

2 ml/l

5 ml/l



Return home

Day after return home

Start of the week

Before basketing



For 1 day  
Demanding flights: for a period of 2-3 days

For a period of 1-2 days

For 2 days

Great speed: 1 day  
Medium distance: 2-3 days



Fast recovery

Proven efficacy in resistant trichomonas

Crop, proventriculus & intestinal health, solid droppings

Form



In warm weather: basketing day

Pinch naris, add drop, allow bird to inhale

Can be administered daily

Before an important race: daily for a period of 10 days in combination with Immuno-Plus



**Need advice?**

Your Neorni veterinarian will be happy to help

►►► This schedule is an indication of how the NeorniPharma products can be used in certain circumstances. We try to develop a complete and appropriate schedule with as few products as possible. All these supplements can be used in combination, if required. In case of poor performance, we will look for the cause with you and decide upon any adjustments or further treatments. Food supplements are best prepared the day before so they can be well absorbed, which will help the pigeons digest the food better.



## Distance – return home



**Immuno Recup Caps**



**Aminovital**



**Trichoryza Drops**



5 ml/l

5 ml/l

5 ml/l or 10 ml/kg  
or 1 drop/naris



For a period of 1-2 days

For a period of 2-3 days

For 2 days



Fast recovery

Fast recovery

Proven efficacy in resistant  
trichomonas

## Distance - intermediate week



**Dysbac Liquid**



**Respicor**



**Hepa-Plus**

61



2 ml/l

3 ml/l

4 ml/l



For 2 days

For 2 days

For 2 days



Crop, proventriculus & intestinal  
health, solid droppings

Respiratory tract, white noses

Liver support

## Distance - before basketing



**Feather Energy**



**Vita Energy**



**Immuno-Plus**



5ml/kg

10g/kg feed

2 ml/l



For a period of 3-4 days

For a period of 3-4 days

For a period of 3-4 days



Promotes recovery

Stimulates fitness

Extra energy



# Moulting

Preparations for good breeding start already during the moulting period. It is important that pigeons are given rest during this period, and that they are supporting by way of moulting vitamins and amino acids such as **Aminovital** and **Condi-Plus**.



<b>Aminovital</b>
2 ml/l
During moulting
2-3 days/week
<b>AMINO ACIDS + VITAMINS</b>

<b>Condi-Plus</b>
2 ml/l
During moulting
2-3 days/week
<b>LYSINE + PLANT EXTRACTS</b>



## Controlling ectoparasites during moulting

Ectoparasites are best checked during moulting and addressed if necessary with **Exmite Spray**, **V-Exmite Drops** or **Vermi-Complete** (in the bath water).

# Preparation for breeding

**62** Once the pigeon season finishes at the end of September/beginning of October, we strongly recommend you **collect mixed manure from the breeders for a period of 5 days and have it tested**. This mixed manure is checked for worms, coccidiosis, yeasts/fungi, and then a culture is grown to check for paratyphoid/salmonella. The results can be used to draw up a vaccination plan.

	Paratyphoid/salmonella <b>NEGATIVE</b>	Paratyphoid/salmonella <b>POSITIVE</b>
<b>Treatment</b>	Not required	Required for a period of 10 days according to antibiogram
<b>Vaccination</b>	<p><b>Strongly recommended</b></p> <ul style="list-style-type: none"> <li>Day 1: salmonella/paratyphoid autovaccine</li> <li>Day 10: paramyxo vaccination (to be repeated annually; can also be done at a different time)</li> <li>Day 21-28: salmonella/paratyphoid autovaccine</li> </ul>	<p><b>Required</b></p> <ul style="list-style-type: none"> <li>2 days after treatment: salmonella/paratyphoid autovaccine</li> <li>10 days later: paramyxo vaccination (to be repeated annually; can also be done at a different time)</li> <li>3-4 weeks later: salmonella/paratyphoid autovaccine</li> </ul>
<b>Time of vaccination</b>	1st vaccination 6-7 weeks prior to mating, 2nd vaccination 4 weeks prior to mating	
<b>Support after vaccination</b>	<b>Immuno-Plus 2 ml/l</b> for a period of 3-5 days	




### Need advice?

Your Neorni veterinarian will be happy to help

### Practical

- Gather manure every day for 5 days and refrigerate it.
- Send the manure in, or take your sample to your veterinarian
- If required, the pigeons can be checked at the same time
- This is not harmful to moulting pigeons





# SCHEDULES FOR PARROTS





# Parrots & parakeets

## Veterinarian check-up

	Private		Professional (breeders)
Accommodation	<i>indoor</i>	<i>outdoor</i>	<i>indoor/outdoor</i>
General check-up + manure tests = nails/bill filing	<ul style="list-style-type: none"> <li>At time of purchase</li> <li>once a year</li> </ul>		<ul style="list-style-type: none"> <li>At time of purchase</li> <li>6-8 weeks prior to breeding</li> </ul>
Extra manure tests	/	Every 6 months	Every 6 months
Worming	If required: <b>Vermi-Complete</b>		



## Support: general & moulting



▶▶▶▶ During moulting, give regular baths/showers/sprays to support plumage. After the moulting period, check your birds for ectoparasites and if needed, treat with **Exmite Spray**.

64

GENERAL VITAMINS	MOULTING VITAMINS	
<b>Vita-Plus / Vita Energy</b>	<b>Aminovital</b>	<b>Condi-Plus</b>
0.2 ml/100 ml water or 1 g/100 g egg food/feed	0.2 ml/100 ml water	0.2 ml/100 ml water
Throughout the year	During moulting	During moulting
1-2 days/week	3 days/week	3 days/week
<b>EXTRA VITAMINS A, B, C, D, E &amp; K</b>	<b>AMINO ACIDS &amp; VITAMINS</b>	<b>PLANT EXTRACTS &amp; LYSINE</b>

## Breeding: preparation & young



▶▶▶▶ Due to the many viral infections possible in psittaciformes, immunity support is the most important aspect when breeding young birds.

PREPARATION	BREEDING YOUNG BIRDS	
<b>Immuno-Plus</b>	<b>Immuno Booster NEW</b>	<b>Immuno-Plus</b>
0.2 ml/100 ml water	Natural brooding: 2 g/100 g egg food/feed Hand rearing: 1-2 g in the mush (spread over several mush feeds)	Natural brooding: 0.2 ml/100 ml water Hand rearing: 0.5-1 ml in the mush or directly into the bill (once a day)
6 weeks prior to mating until the 1st clutch	Daily from birth until independence	Daily from birth until independence
Daily	Daily	Daily
<b>STIMULATES SPERM CELL &amp; EGG CELL PRODUCTION</b>	<b>COMPLETE PREMIX FOR GROWING BIRDS &amp; IMMUNITY SUPPORT</b>	<b>EXTRA IMMUNITY SUPPORT FOR YOUNG BIRDS</b>

# Index

Aminovital .....	42
Condi-Plus .....	43
CU-For Liquid .....	38
Derma-Plus .....	28
Dysbacteriose .....	18
Dysbac Liquid .....	19
Exmite Spray .....	29
Feather Energy .....	23
Freeson Drops .....	22
Hepa-Plus .....	14
Immuno Booster NEW .....	7
Immuno-Plus .....	8
Immuno Recup Caps .....	9
Natural breeding promotion .....	6
Neorni Stabiliser .....	33
PikStop .....	24
Probi-Plus .....	34
Ready .....	15
Respicor .....	25
SilverFlox .....	39
SilverSpray .....	35
Skin Spray .....	36
Super Start-Up .....	10
Trichoryza Drops .....	40
Vita Energy .....	44
Vita-Min-DCP .....	45
Vita-Plus .....	46
Vit E + Se .....	11
Schedules for pigeons .....	57
Breeding .....	58
Moulting .....	62
Racing season .....	60-61
Flight preparation.....	59
Schedules for parrots .....	63
Parrots & parakeets .....	64
Schedules for songbirds .....	47
Annual schedule for Euro & Canaries .....	49-50
Annual schedule for Euro & Canaries Timeline .....	48
Annual schedule for Exotics .....	54-55
Breeding: weaned young .....	53
Breeding: hatching to weaning .....	52
Breeding preparation .....	51

# Export prices

Please find the NeorniPharma export prices for private individuals and professionals below. These prices are **exclusive of VAT**. The VAT rates for the country of destination are applied when shipping. As always, professionals are VAT exempt.

PRODUCT	€ excl. VAT	PRODUCT	€ excl. VAT
Immuno Booster NEW (1 kg)	53,72	CU-For Liquid (250 ml)	12,40
Immuno-Plus (100 ml)	20,66	SilverFlox (1 l)	26,45
Immuno-Plus (250 ml)	45,45	Trichoryza Drops (50 ml)	14,05
Immuno-Plus (1 l)	147,93	Trichoryza Drops (250 ml)	48,76
Immuno Recup Caps (80 c)	19,83	Aminovital (100 ml)	15,70
Super Start-Up (200 g)	20,66	Aminovital (250 ml)	28,93
Vit E + Se (100 ml)	14,05	Aminovital (1 l)	70,25
Vit E + Se (250 ml)	28,93	Condi-Plus (100 ml)	12,40
Hepa-Plus (100 ml)	16,53	Condi-Plus (250 ml)	21,49
Hepa-Plus (250 ml)	29,75	Condi-Plus (1 l)	68,60
Hepa-Plus (1 l)	79,34	Vita Energy (250 g)	20,66
Ready (100 ml)	18,18	Vita Energy (1 kg)	61,98
Ready (250 ml)	39,67	Vita-Min-DCP (100 ml)	14,05
Ready (1 l)	119,01	Vita-Min-DCP (250 ml)	26,45
Dysbac Liquid (250 ml)	30,58	Vita-Min-DCP (1 l)	72,73
Freeson Drops (30 ml)	14,05	Vita-Plus (100 ml)	14,88
Feather Energy (400 ml)	23,97	Vita-Plus (250 ml)	23,14
PikStop (100 ml)	23,14	Vita-Plus (1 l)	68,60
Respicor (100 ml)	15,70	PURE	
Respicor (250 ml)	28,93	Brewer's yeast (1 kg)	14,88
Derma-Plus (50 ml)	28,93	Dextrose (1 kg)	14,88
Exmite Spray (250 ml)	18,18	Elektrolyt (500 g)	14,88
Exmite Spray (1 l)	57,02	Yellow (100 g)	12,40
Neorni Stabilizer (1 l)	16,53	Yellow (500 g)	49,59
Probi-Plus (200 g)	14,88	Highly digestible protein (400 g)	20,66
SilverSpray (500 ml)	12,40	Red (100 g)	23,14
Skin Spray (200 ml)	14,05	Red (500 g)	70,25

# Dosage chart (part 1)

## NEORNI

Beverloosesteenweg 129, Paal  
 ••• Tel. +32 (0)11 96 91 02...•••



Veterinary practice



Laboratory



Product range



## CAPSULES

- Alternative administration of capsules via drinking water (ml) or feed (g)
- For all cases where inserting is not possible

Capsule	/100 ml	/50 g
SBS Caps	1	1
Asper Dox Caps	1	1
Endo-Ecto Plus Caps 0.5 kg	-	1
Orni Caps	-	1
Sulfaclozine Caps	1	1-2
<b>Capsule</b>	<b>/300 ml</b>	<b>/150 g</b>
Endo-Ecto Plus Caps 1 kg	-	1
Tricho Cox Caps	1	1



**Neornipharma**  
 powders & capsules

# Dosage chart (part 2)

## POWDERS



- Easy administration of powders via drinking water (l) or feed (g/kg), without using a scale
- Level off\* the spoon after scooping

Powder 'PURE'	
Amoxicillin 80%	1 / 1
Colistin 480IU/mg	2 / 1
Doxycycline 15%	2 / 1
Doxycycline 50%	1 / 21
Lincospectin 66%	1 / 1,51
Spiramycin 50%	1 / 1,51
Tiamulin 45%	1 / 21
Trim-Sulfa 90%	1 / 1,51
Tylosin 100%	2 / 1

\*Spoon included with your purchase



Powder	
Aitree	4 / 1
Bronchifree	3 / 1
CAS (loft disinfection)	3 / 1
Coli-Mix	3 / 1
Gould Mix	3 / 1
Immuno Booster NEW	2 / 100 g
Megadysbac	3 / 1
Mucolyt	3 / 1
Probi-Plus	2 / 1
Red Race	3 / 1
Sulfaclozine	2 / 1
Super Start-Up	4 / kg
Tricho Geel	3 / 21 (<0.5kg 1 / 1)
Vita Energy	5 / kg
YPD Mix	3 / 1



**Visit us**  
 NeorinPharma  
 Beverlosesteenweg 129  
 3583 Paal, Belgium

**Contact us**  
 T +32 (0)11 96 91 02  
 E info@neorinpharma.be  
 W www.neorinpharma.be

**Download our catalogues**



**Follow us on Facebook**



**NeorinPharma, custom formulations for your birds**  
 The dosages on this card offer an alternative administration compared to the product label, tailored to your bird or situation. Contact us if in doubt.



EN

T +32 (0)11 96 91 02  
E [info@neornipharma.be](mailto:info@neornipharma.be)  
W [www.neornipharma.be](http://www.neornipharma.be)

Desk & depot:

Beverlosesteenweg 129,  
3583 Paal, Belgium

